

New Test

3000479

Criteria Systemic Sclerosis Panel

SSC PANEL



Additional Technical Information

Methodology: Semi-Quantitative Indirect Fluorescent Antibody/Semi-Quantitative Multiplex Bead Assay/Semi-Quantitative Enzyme-Linked Immunosorbent Assay
Performed: Sun, Tue, Thu
Reported: 1-3 days

Specimen Required: Collect: Serum Separator Tube (SST).
Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 3 mL serum to an ARUP Standard Transport Tube. (Min: 0.25 mL)
Storage/Transport Temperature: Refrigerated.
Unacceptable Conditions: Contaminated, hemolyzed, or severely lipemic specimens.
Stability (collection to initiation of testing): Ambient: 48 hours; Refrigerated: 2 weeks; Frozen: 1 year (avoid repeated freeze/thaw cycles)

Reference Interval:

Test Number	Components	Reference Interval	
3000082	Antinuclear Antibody (ANA), HEp-2, IgG	Less than 1:80	
0050599	Scleroderma (Scl-70) (ENA) Antibody, IgG	29 AU/mL or less	Negative
		30-40 AU/mL	Equivocal
		41 AU/mL or greater	Positive
2001601	RNA Polymerase III Antibody, IgG	19 Units or less	Negative
		20-39 Units	Weak Positive
		40-80 Units	Moderate Positive
		81 Units or greater	Strong Positive

Interpretive Data: Refer to report.

CPT Code(s): 86039; 86235; 83516

New York DOH Approved.

HOTLINE NOTE: Refer to the Test Mix Addendum for interface build information.