

Patient: [REDACTED]
DOB: [REDACTED] **Age:** [REDACTED] **Gender:** [REDACTED]
Patient Identifiers: [REDACTED]
Visit Number (FIN): [REDACTED]

Client: [REDACTED]
Physician: [REDACTED]

ARUP Test Code: 0081062
 Collection Date: 11/23/2016
 Received in lab: 11/30/2016
 Completion Date: 12/05/2016

Patient Information Used in Risk Calculations:

Maternal Age at Delivery: [REDACTED] yr
 Estimated Due Date: 05/27/2017
 Gestational Age at Draw: 13 Weeks 4 Day(s)
 Maternal Weight: 212 lbs
 Maternal Race: [REDACTED]
 Number of Fetuses: One
 Family History of Aneuploidy: No
 Crown Rump Length: 5.77 cm

Marker Measurement MoM

PAPP-A 1732 mIU/L 1.00
 NT Not Done

Sonographer Name: N/A
 Sonographer Cert #: N/A
 Ultrasound Date: 11/14/2016

Interpretation: INTERPRETATION PENDING 2nd SPECIMEN

The ideal time to draw the 2nd sample for the Integrated screen is between 16 and 18 weeks gestation. For this patient, ideally collect between 10-Dec-16 and 24-Dec-16.

However, the sample may be drawn as early as 3-Dec-16 or as late as 10-Feb-17 and still provide valid information.

The PAPP-A test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS



Patient: [REDACTED]
 ARUP Accession: 16-328-135453