

Ankylosing Spondylitis (HLA-B27) Genotyping

Ankylosing spondylitis (AS) is a chronic form of arthritis in which the sacroiliac joints become inflamed, leading to back pain and, in some cases, the formation of bone between vertebrae (ie, ankylosis). Although spinal involvement is typical to AS, other areas of the body (eg, shoulders, hips, knees, etc.) may also become stiff or inflamed. Although no specific cause of AS has yet been identified, data support a strong correlation between AS and the presence of HLA-B27; 80-90% of AS patients are HLA-B27 positive, compared to 5-10% of the general population.¹

Genetics

Gene

HLA-B27 group alleles

Allele(s)

More than 213 known alleles²

Inheritance

Codominant

Prevalence

Overall prevalence in North America is 6-8% but varies by ethnicity. Notably, there is a descending gradient from north to south of the HLA-B27 allelic prevalence; in particular, the ancestral HLA-B*27:05 allele frequency ranges from >20% in Northern Europe to <1% in sub-Saharan Africa.

Test Interpretation

Sensitivity/Specificity

- Analytic sensitivity/specificity: >99%
- Clinical sensitivity: 90%
- Clinical specificity: <1% in unaffected individuals with no family history of AS

Results

Result	Interpretive Note
Positive: HLA-B27 was detected	Result supports a clinical diagnosis of AS or another related disorder (eg, Reiter syndrome, anterior uveitis, psoriatic arthritis, and inflammatory bowel disease)
Negative: HLA-B27 was not detected	Does not rule out AS; refer to clinical sensitivity

Limitations

- · Contraindicated for prenatal or carrier testing
- · Rare diagnostic errors can occur due to primer site variants.
- This test does not rule out alleles HLA-B*27:06 or HLA-B*27:09, which are not associated with spondyloarthropathies.
- Certain rare alleles present in <1% of the population will not be detected.
- · Other rare or uncharacterized alleles are possible and may lead to false positive or false negative results.

Featured ARUP Testing

Ankylosing Spondylitis (HLA-B27) Genotyping 0050392

Method: Polymerase Chain Reaction (PCR)/Fluorescence Monitoring

- Not a diagnostic test for AS
- May assist in the diagnosis of AS only if other clinical signs and symptoms are present
- Test should not be performed for prenatal diagnosis of AS because a positive result is not predictive for the disorder

References

- 1. Wenker KJ, Quint JM. Ankylosing spondylitis. In: StatPearls. StatPearls Publishing; 2022. [Updated: Apr 2022; Accessed: Aug 2022]
- 2. Khan MA. An update on the genetic polymorphism of HLA-B*27 with 213 alleles encompassing 160 subtypes (and still counting). Curr Rheumatol Rep. 2017;19(2):9.
- 3. Profaizer T, Dibb K, Bethers H, et al. Comparison of next-generation sequencing-based human leukocyte antigen typing with clinical flow cytometry and allele-specific PCR melting assays for HLA-B27 genotyping. *J Appl Lab Med*. 2021;6(5):1221-1227.

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology. 500 Chipeta Way, Salt Lake City, UT 84108 (800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com
Content Review August 2022 | Last Update June 2023

© 2023 ARUP Laboratories. All Rights Reserved.

Client Services - (800) 522-2787