

Polymyositis Panel

Last Literature Review: March 2024 Last Update: January 2025

Myositis is characterized by inflammation of the skeletal muscles involved in movement.^{1,2} The detection of antibodies may help to establish a diagnosis, aid in prognosis, and support treatment decisions.

Disease Overview

Myositis may occur in a number of inflammatory myopathies, including polymyositis/antisynthetase syndrome, dermatomyositis, necrotizing autoimmune myopathy, and sporadic inclusion body myositis, as well as overlap syndromes with [connective tissue diseases](#).^{1,2} The primary symptom of all forms of myositis is progressive muscle weakness that may develop over a period of weeks, months, or years.^{1,2} Other symptoms may include joint pain and fatigue.^{1,2}

Antibody testing for myositis should be considered after a standard workup for inflammatory myopathies because it may aid in distinguishing between myopathies,^{1,2} which can have important implications for therapy and prognosis.

Refer to the ARUP Consult [Inflammatory Myopathies – Myositis](#) topic for more information about myositis and the typical testing strategy for inflammatory myopathies.

Test Description

This antibody panel test may be useful for the evaluation of patients with progressive proximal muscle weakness and antisynthetase syndrome. Clinical phenotypes for specific antibody-associated inflammatory myopathies often overlap, and targeted panels allow for rapid identification of associated antibodies. Use of the most targeted panel, ie, the panel that most closely matches the patient's complete clinical phenotype, is recommended.

Featured ARUP Testing

Polymyositis Panel 3018868

Method: Qualitative Immunoprecipitation / Semi-Quantitative Multiplex Bead Assay / Semi-Quantitative Indirect Fluorescent Antibody (IFA) / Qualitative Immunoblot

Additional ARUP Myositis Panels

ARUP Panel to Consider	Clinical Utility	Additional Test Information
Dermatomyositis and Polymyositis Panel 3018866 Antibodies overlap with the antibodies on this panel	May be useful for the evaluation of patients with progressive proximal muscle weakness and/or with cutaneous manifestations suggestive of dermatomyositis	Dermatomyositis and Polymyositis Panel Test Fact Sheet
Dermatomyositis Autoantibody Panel 3018870 Includes antibodies that are specific to dermatomyositis	May be useful for the evaluation of patients with characteristic cutaneous manifestations of dermatomyositis with or without muscle weakness	Dermatomyositis Autoantibody Panel Test Fact Sheet
Extended Myositis Panel 3018867 Antibodies overlap with the antibodies on this panel	May be useful for the evaluation of patients with progressive proximal muscle weakness and/or other clinical findings suggestive of polymyositis/antisynthetase syndrome, dermatomyositis, necrotizing autoimmune myopathy, or overlap syndromes associated with connective tissue disease	Extended Myositis Panel Test Fact Sheet

ARUP Panel to Consider	Clinical Utility	Additional Test Information
Interstitial Lung Disease Autoantibody Panel 3018869 Antibodies overlap with the antibodies on this panel	May be useful for the evaluation of patients with interstitial lung disease with or without other signs and symptoms of myositis	Interstitial Lung Disease Autoantibody Panel Test Fact Sheet

Antibodies Tested

This panel detects a selection of antibodies specific to or associated with myositis. For more information about the clinical associations with each of these antibodies, visit the ARUP Consult [Inflammatory Myopathies – Myositis](#) topic.

Polymyositis Panel: Antibodies Detected and Methodology	
Myositis-Specific Antibodies ^a	
Antibody	Method
EJ (glycyl-tRNA synthetase) Ab	Qualitative immunoprecipitation
Ha (tyrosyl-tRNA synthetase) Ab	Qualitative immunoblot and qualitative immunoprecipitation
Ks (asparaginyl-tRNA synthetase) Ab	Qualitative immunoblot and qualitative immunoprecipitation
Jo-1 (histidyl-tRNA synthetase) Ab, IgG	Semiquantitative multiplex bead assay
OJ (isoleucyl tRNA synthetase) Ab	Qualitative immunoprecipitation
PL-7 (threonyl-tRNA synthetase) Ab	Qualitative immunoprecipitation
PL-12 (alanyl-tRNA synthetase) Ab	Qualitative immunoprecipitation
SRP (signal recognition particle) Ab	Qualitative immunoprecipitation
Zo (phenylalanyl-tRNA synthetase) Ab	Qualitative immunoblot and qualitative immunoprecipitation
Myositis-Associated Antibodies	
Antibody	Method
Antinuclear Ab (ANA), Hep-2, IgG ^b	Semiquantitative indirect fluorescent antibody

^aMyositis-specific antibodies are generally regarded as mutually exclusive with rare exceptions. The occurrence of two or more myositis-specific antibodies should be carefully evaluated in the context of the patient's clinical presentation. Refer to the ARUP Consult [Inflammatory Myopathies – Myositis](#) topic for more information about myositis.

^bThe presence of ANA is a feature of systemic autoimmune rheumatic diseases, however, ANA lacks diagnostic specificity and may occur in the general population. Positive ANA must be confirmed by more specific serologic tests. For more information, refer to the [Antinuclear Antibody \(ANA\) With Hep-2 Substrate Test Fact Sheet](#).

Ab, antibody; IgG, immunoglobulin G

Some antibodies may be orderable separately; refer to the [ARUP Laboratory Test Directory](#).

Test Interpretation

Results

- **Positive:** Antibody detected.

- Supports a clinical diagnosis of polymyositis/antisynthetase syndrome, necrotizing autoimmune myopathy, and/or an overlap syndrome.
- Results for specific antibodies may be reported as low/weak positive, positive, or high/strong positive.
- Additional interpretive information for positive antibodies may be provided on the Patient Report.
- Myositis-specific antibodies are generally regarded as mutually exclusive with rare exceptions; the occurrence of two or more myositis-specific antibodies should be carefully evaluated in the context of the patient's clinical presentation.
- Myositis-associated antibodies may be found in patients with overlap syndromes and other conditions and are generally not specific for myositis.
- **Negative:** Antibody not detected.

Limitations

Results are not diagnostic in the absence of other findings and should be considered in the complete clinical context.

Negative results do not rule out a diagnosis of inflammatory myopathy or overlap syndrome.

References

1. Selva-O'Callaghan A, Pinal-Fernandez I, Trallero-Araguás E, et al. [Classification and management of adult inflammatory myopathies](#). *Lancet Neurol*. 2018;17(9):816-828.
2. Schmidt J. [Current classification and management of inflammatory myopathies](#). *J Neuromuscul Dis*. 2018;5(2):109-129.

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology. 500 Chipeta Way, Salt Lake City, UT 84108
(800) 522-2787 | (801) 583-2787 | aruplab.com | arupconsult.com