

# Noonan Spectrum Disorders Panel

Noonan spectrum disorders (NSDs) are a group of genetic syndromes caused by pathogenic germline variants in genes in the Ras/mitogen activated protein kinase (MAPK) pathway, which controls the cell cycle and cell differentiation. The vast majority of causative variants increase pathway signaling; thus, the resulting syndromes exhibit phenotypic overlap and share a predisposition for developing malignancies.

## Disease Overview

### Symptoms of Noonan Syndrome (NS)

- Characteristic facial features
- Short stature
- Broad webbed neck (fetal cystic hygroma/increased nuchal translucency)
- Congenital heart defect
- Developmental delay
- Undescended testes
- Coagulation defects
- Lymphatic dysplasias

### Etiology of NSDs

Pathogenic sequence variants in Ras pathway genes

### Prevalence

NS: 1/1,000-2,500

### Inheritance

Autosomal dominant for all analyzed genes

### Genotype-Phenotype Correlation

Variants in multiple genes cause overlapping phenotypes for NSD

## Test Description

See [Genes Tested](#) table for genes included in this panel.

## Clinical Sensitivity

Dependent on clinical phenotype

- Approximately 99% for cardiofaciocutaneous syndrome (CFCS)<sup>1</sup>
- Approximately 80-90% for Costello syndrome (CS)<sup>2,3,4,5</sup>
- Approximately 70-80% for NS<sup>6-13</sup>

## Featured ARUP Testing

### Noonan Spectrum Disorders Panel, Sequencing 2010772

**Method:** Massively Parallel Sequencing

Confirm a suspected clinical diagnosis of:

- Noonan syndrome (NS)
- Cardiofacial cutaneous syndrome (CFCS)
- Costello syndrome (CS)
- Legius syndrome (LS)
- Noonan syndrome with multiple lentigines (LEOPARD syndrome)
- Noonan-like syndrome
- Noonan-like syndrome with loose anagen hair (NS/LAH)

Given the genotypic and phenotypic overlap among NSDs, the NSD panel is the recommended first-line test for determining a genetic etiology.

**Contraindications:** This panel should not be ordered in individuals with primary juvenile myelomonocytic leukemia (JMML) as the assay may not detect mosaicism for somatic variants associated with malignancy.

### Noonan Spectrum Disorders Panel, Sequencing, Fetal 2010769

**Method:** Massively Parallel Sequencing

Confirm a diagnosis of an NSD in a pregnancy with clinically suggestive findings, such as increased nuchal translucency, cystic hygroma, and cardiac defects. For a fetus with ultrasonographic abnormalities, genomic microarray should be ordered prior to the NSD panel.

If a familial sequence variant has been previously identified, targeted sequencing for that variant may be appropriate; refer to the [Laboratory Test Directory](#) for additional information.

## Limitations

- A negative result does not exclude a diagnosis of a MAPK pathway disorder.
- Diagnostic errors can occur due to rare sequence variations.
- Interpretation of this test result may be impacted if the individual has had an allogeneic stem cell transplantation.
- The following will not be evaluated:
  - Variants outside the coding regions and intron-exon boundaries of the targeted genes
  - Regulatory region variants and deep intronic variants
  - Large deletions/duplications
  - Noncoding transcripts
- The following may not be detected:
  - Deletions/duplications/insertions of any size by massively parallel sequencing
  - Some variants due to technical limitations in the presence of pseudogenes, repetitive, or homologous regions
  - Low-level somatic variants

## Analytic Sensitivity

For massively parallel sequencing:

Variant Class	Analytic Sensitivity (PPA) Estimate <sup>a</sup> (%)	Analytic Sensitivity (PPA) 95% Credibility Region <sup>a</sup> (%)
SNVs	99.2	96.9-99.4
Deletions 1-10 bp	93.8	84.3-98.2
Deletions 11-44 bp	100	87.8-100
Insertions 1-10 bp	94.8	86.8-98.5
Insertions 11-23 bp	100	62.1-100

bp, base pairs; PPA, positive percent agreement; SNVs, single nucleotide variants

## Genes Tested

Gene	Alias Symbol(s)	MIM Number	Disorder
<b><i>BRAF</i></b>	BRAF1	164757	CFCS 1 NS 1 NS 7 LEOPARD syndrome 3
<b><i>CBL</i></b>	CBL2, RNF55, c-Cbl	165360	NS-like disorder with or without juvenile myelomonocytic
<b><i>HRAS</i></b>	HRAS1	190020	Melanocytic Nevus syndrome, congenital Schimmelpenning-Feuerstein-Mims syndrome CS
<b><i>KRAS</i></b>	KRAS2, KRAS1	190070	Schimmelpenning-Feuerstein-Mims syndrome NS 3 CFCS 2
<b><i>LZTR1</i></b>	LZTR-1, BTBD29	600574	NS 10

Gene	Alias Symbol(s)	MIM Number	Disorder
<b>MAP2K1</b>	PRKMK1, MEK1, MAPKK1	176872	NS 1 CFCS 3
<b>MAP2K2</b>	PRKMK2, MEK2	601263	CFCS 4
<b>NRAS</b>	N-ras	164790	Schimmelpenning-Feuerstein-Mims syndrome NS 6
<b>PTPN11</b>	NS1, BPTP3, SH-PTP2, SHP-2, PTP2C, SHP2	176876	LEOPARD syndrome 1 NS 1
<b>RAF1</b>	Raf-1, c-Raf, CRAF	164760	NS 5 LEOPARD syndrome 2
<b>RASA2</b>	GAP1M	601589	
<b>RIT1</b>	RIT, RIBB, ROC1, MGC125864, MGC125865	609591	NS 8
<b>SHOC2</b>	KIAA0862, SOC2, SUR-8, SOC-2, SUR8	602775	NS-like disorder with loose anagen hair 1
<b>SOS1</b>	GINGF, HGF, GF1	182530	NS 4
<b>SOS2</b>		601247	NS 9
<b>SPRED1</b>	FLJ33903, PPP1R147	609291	Legius syndrome

## References

1. Rauen KA. [The RASopathies](#). *Annu Rev Genomics Hum Genet*. 2013;14:355-369.
2. Aoki Y, Niihori T, Kawame H, et al. [Germline mutations in HRAS proto-oncogene cause Costello syndrome](#). *Nat Genet*. 2005;37(10):1038-1040.
3. Estep AL, Tidyman WE, Teitell MA, et al. [HRAS mutations in Costello syndrome: detection of constitutional activating mutations in codon 12 and 13 and loss of wild-type allele in malignancy](#). *Am J Med Genet A*. 2006;140(1):8-16.
4. Gripp KW, Lin AE, Stabley DL, et al. [HRAS mutation analysis in Costello syndrome: genotype and phenotype correlation](#). *Am J Med Genet A*. 2006;140(1):1-7.
5. Kerr B, Delrue MA, Sigaudy S, et al. [Genotype-phenotype correlation in Costello syndrome: HRAS mutation analysis in 43 cases](#). *J Med Genet*. 2006;43(5):401-405.
6. Tartaglia M, Kalidas K, Shaw A, et al. [PTPN11 mutations in Noonan syndrome: molecular spectrum, genotype-phenotype correlation, and phenotypic heterogeneity](#). *Am J Hum Genet*. 2002;70(6):1555-1563.
7. Roberts AE, Araki T, Swanson KD, et al. [Germline gain-of-function mutations in SOS1 cause Noonan syndrome](#). *Nat Genet*. 2007;39(1):70-74.
8. Tartaglia M, Pennacchio LA, Zhao C, et al. [Gain-of-function SOS1 mutations cause a distinctive form of Noonan syndrome](#). *Nat Genet*. 2007;39(1):75-79.
9. Aoki Y, Niihori T, Inoue Sichi, et al. [Recent advances in RASopathies](#). *J Hum Genet*. 2016;61(1):33-39.
10. Schubbert S, Zenker M, Rowe SL, et al. [Germline KRAS mutations cause Noonan syndrome](#). *Nat Genet*. 2006;38(3):331-336.
11. Brasil ASalem, Pereira AC, Wanderley LTurolla, et al. [PTPN11 and KRAS gene analysis in patients with Noonan and Noonan-like syndromes](#). *Genet Test Mol Biomarkers*. 2010;14(3):425-432.
12. Sarkozy A, Carta C, Moretti S, et al. [Germline BRAF mutations in Noonan, LEOPARD, and cardiofaciocutaneous syndromes: molecular diversity and associated](#)

[phenotypic spectrum](#). *Hum Mutat*. 2009;30(4):695-702.

13. Nava C, Hanna N, Michot C, et al. [Cardio-facio-cutaneous and Noonan syndromes due to mutations in the RAS/MAPK signalling pathway: genotype-phenotype relationships and overlap with Costello syndrome](#). *J Med Genet*. 2007;44(12):763-771.

## Related Information

[Laboratory Testing for Developmental Delay, Intellectual Disability, and Autism Spectrum Disorder](#)  
[Testing for Genetic Syndromes Related to Developmental Delay \(DD\), Intellectual Disability \(ID\), and Autism Spectrum Disorder \(ASD\)](#)

ARUP Laboratories is a nonprofit enterprise of the University of Utah and its Department of Pathology, 500 Chipeta Way, Salt Lake City, UT 84108  
(800) 522-2787 | (801) 583-2787 | [aruplab.com](#) | [arupconsult.com](#)  
Content Review February 2019 | Last Update October 2023