

Patient Name \_\_\_\_\_ Date of Birth \_\_\_\_\_

Ordering Physician \_\_\_\_\_ Physician's Office Contact \_\_\_\_\_

Office Phone \_\_\_\_\_

## Patient History for Prolonged Clot Time Reflex Panel

This panel is designed to evaluate prolonged clotting times when cause is unknown. Condition-specific Testing is recommended if the cause for the prolonged clotting time is known (e.g., factor deficiency, inhibitor, or von Willebrand disease).

Identify the clinical presentation related to the work-up of a prolonged clotting time.

- Bleeding
- Thrombosis
- Unexpected clotting time prolongation

Which clotting time test is prolonged at the referring location?

- PT
- aPTT
- dRVVT
- Thrombin Time

If patient has taken any coagulation-related medication in the past 7 days, indicate date last given \_\_\_\_\_

Current anticoagulant medications may interfere with the testing and cause erroneous results.

- Coumadin (warfarin)
- Unfractionated heparin
- Arixtra (fondaparinux)
- Vitamin K
- Thrombolytic (such as t-PA)
- Direct thrombin inhibitor: Pradaxa (dabigatran), Acova (argatroban), Angiomax (bilvalirudin)
- Low-molecular-weight heparin: Lovenox (enoxaparin), Fragmin (dalteparin)
- Direct Xa inhibitor: Xarelto (rivaroxaban), Eliquis (apixaban), Savaysa (edoxaban)

Has transfusion or replacement factor been given within the past 72 hours? Treatment may affect testing results.

Yes  No

- DDAVP
- Cryoprecipitate
- Fresh frozen plasma
- VWF concentrate
- FVIII concentrate
- IX concentrate
- Other product; specify \_\_\_\_\_

---

Email completed form to Coagulation811@aruplab.com

---