

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 4/3/1979
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Lead, Blood (Venous)

ARUP test code 0020098

Lead, Blood (Venous)

<2.0 ug/dL (Ref Interval: <=4.9)

INTERPRETIVE INFORMATION: Lead, Blood (Venous)

Analysis performed by Inductively Coupled Plasma-Mass Spectrometry (ICP-MS).

Elevated results may be due to skin or collection-related contamination, including the use of a noncertified lead-free tube. If contamination concerns exist due to elevated levels of blood lead, confirmation with a second specimen collected in a certified lead-free tube is recommended.

Information sources for blood lead reference intervals and interpretive comments include the CDC's "Childhood Lead Poisoning Prevention: Recommended Actions Based on Blood Lead Level" and the "Adult Blood Lead Epidemiology and Surveillance: Reference Blood Lead Levels (BLLs) for Adults in the U.S." Thresholds and time intervals for retesting, medical evaluation, and response vary by state and regulatory body. Contact your State Department of Health and/or applicable regulatory agency for specific guidance on medical management recommendations.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

Group	Concentration	Comment
Children	3.5-19.9 ug/dL	Children under the age of 6 years are the most vulnerable to the harmful effects of lead exposure. Environmental investigation and exposure history to identify potential sources of lead. Biological and nutritional monitoring are recommended. Follow-up blood lead monitoring is recommended.
	20-44.9 ug/dL	Lead hazard reduction and prompt medical evaluation are recommended. Contact a Pediatric Environmental Health Specialty unit or poison control center for guidance.

H=High, L=Low, *=Abnormal, C=Critical

	Greater than 44.9 ug/dL	Critical. Immediate medical evaluation, including detailed neurological exam is recommended. Consider chelation therapy when symptoms of lead toxicity are present. Contact a Pediatric Environmental Health Specialty Unit or poison control center for assistance.
Adult	5-19.9 ug/dL	Medical removal is recommended for pregnant women or those who are trying or may become pregnant. Adverse health effects are possible. Reduced lead exposure and increased blood lead monitoring are recommended.
	20-69.9 ug/dL	Adverse health effects are indicated. Medical removal from lead exposure is required by OSHA if blood lead level exceeds 50 ug/dL. Prompt medical evaluation is recommended.
	Greater than 69.9 ug/dL	Critical. Immediate medical evaluation is recommended. Consider chelation therapy when symptoms of lead toxicity are present.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Lead, Blood (Venous)	22-298-103610	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: