

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB 6/21/2018
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Lead, Blood (Capillary)

ARUP test code 0020745

Lead, Blood (Capillary) <2.0 ug/dL (Ref Interval: <=4.9)

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: Lead, Blood (Capillary)

Elevated results may be due to skin or collection-related contamination, including the use of a noncertified lead-free collection/transport tube. If contamination concerns exist due to elevated levels of blood lead, confirmation with a venous specimen collected in a certified lead-free tube is recommended.

Repeat testing is recommended prior to initiating chelation therapy or conducting environmental investigations of potential lead sources. Repeat testing collections should be performed using a venous specimen collected in a certified lead-free collection tube.

Information sources for reference intervals and interpretive comments include the "CDC Response to the 2012 Advisory Committee on Childhood Lead Poisoning Prevention Report" and the "Recommendations for Medical Management of Adult Lead Exposure, Environmental Health Perspectives, 2007." Thresholds and time intervals for retesting, medical evaluation, and response vary by state and regulatory body. Contact your State Department of Health and/or applicable regulatory agency for specific guidance on medical management recommendations.

Age	Concentration	Comment
All ages	5-9.9 ug/dL	Adverse health effects are possible, particularly in children under 6 years of age and pregnant women. Discuss health risks associated with continued lead exposure. For children and women who are or may become pregnant, reduce lead exposure.
All ages	10-19.9 ug/dL	Reduced lead exposure and increased biological monitoring are recommended.
All ages	20-69.9 ug/dL	Removal from lead exposure and prompt medical evaluation are recommended. Consider chelation therapy when concentrations exceed 50 ug/dL and symptoms of lead toxicity are present.
Less than 19 years of age	Greater than 44.9 ug/dL	Critical. Immediate medical evaluation is recommended. Consider chelation therapy when symptoms of lead toxicity are present.
Greater than 19 years of age	Greater than 69.9 ug/dL	Critical. Immediate medical evaluation is recommended. Consider chelation therapy when symptoms of lead toxicity are present.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

H=High, L=Low, *=Abnormal, C=Critical

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Lead, Blood (Capillary)	20-171-112460	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical