

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 7/23/1958  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Procalcitonin**

ARUP test code 0020763

Procalcitonin

<0.07 ng/mL (Ref Interval: 0.00-0.07)

INTERPRETIVE INFORMATION: Procalcitonin

Procalcitonin levels above 2.00 ng/mL on the first day of ICU admission represent a high risk for progression to severe sepsis and/or septic shock.

Procalcitonin < 0.50 ng/mL:  
Procalcitonin levels below 0.50 ng/mL on the first day of ICU admission represent a low risk for progression to severe sepsis and/or septic shock.

If the procalcitonin measurement is performed shortly after the systemic infection process has started (usually less than 6 hours), these values may still be low. As various non-infectious conditions are known to induce procalcitonin as well, procalcitonin levels between 0.5 ng/mL and 2.00 ng/mL should be reviewed carefully to take into account the specific clinical background and condition(s) of the individual patient.

Performed At: U HOSPITAL LAB (ARUP)  
UNIVERSITY HOSPITAL  
CLINICAL LABORATORY  
SALT LAKE CITY, UT 84132  
Medical Director: LAUREN NICHOLE PEARSON, DO  
CLIA Number: 46D0678613

INTERPRETIVE INFORMATION: Procalcitonin

Procalcitonin > 2.00 ng/mL:  
Procalcitonin levels above 2.00 ng/mL on the first day of ICU admission represent a high risk for progression to severe sepsis and/or septic shock.

Procalcitonin < 0.50 ng/mL:  
Procalcitonin levels below 0.50 ng/mL on the first day of ICU admission represent a low risk for progression to severe sepsis and/or septic shock.

If the procalcitonin measurement is performed shortly after the systemic infection process has started (usually less than 6 hours), these values may still be low. As various non-infectious conditions are known to induce procalcitonin as well, procalcitonin levels between 0.5 ng/mL and 2.00 ng/mL should be reviewed carefully to take into account the specific clinical background and condition(s) of the individual patient.

**H=High, L=Low, \*=Abnormal, C=Critical**



VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Procalcitonin	22-316-702273	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical