

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 4/25/1976
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

von Willebrand Multimeric Panel

ARUP test code 0030002

Factor VIII, Activity 71 % (Ref Interval: 56-191)
REFERENCE INTERVAL: Factor VIII, Activity
Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

von Willebrand Factor, Activity (RCF) 124 % (Ref Interval: 51-215)
REFERENCE INTERVAL: von willebrand Factor, Activity (RCF)
Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

von Willebrand Factor, Antigen 127 % (Ref Interval: 52-214)
REFERENCE INTERVAL: von willebrand Factor, Antigen
Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

von Willebrand Multimeric See Note
von willebrand factor multimeric analysis shows a normal multimeric distribution.
A normal von willebrand panel combined with a normal multimeric distribution does not rule out von willebrand disease since von willebrand values can fluctuate over time. If clinically indicated, consider repeating the von willebrand panel. Additional information regarding diagnosis and subtyping of von willebrand disease is available at www.arupconsult.com.
von willebrand factor and factor VIII are acute phase reactants. Increased values are seen in pregnancy, with estrogen-containing medications, inflammation, infection, liver disease, malignancy, and with exercise, stress, or trauma, including traumatic venipuncture. Acute phase reactions can mask low baseline values in patients with von willebrand disease.
Lower baseline von willebrand and factor VIII values are found in blood type O individuals. Decreased values for von willebrand factor and/or factor VIII activity may also be due to improper sample handling.

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: Von willebrand Multimeric

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Factor VIII, Activity	22-334-149156	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
von Willebrand Factor, Activity (RCF)	22-334-149156	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
von Willebrand Factor, Antigen	22-334-149156	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
von Willebrand Multimeric	22-334-149156	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: