

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 9/6/1959  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Factor XII, Activity**

ARUP test code 0030115

Factor XII, Activity

See Note %

(Ref Interval: 58-166)

Factor XII activity is at least 136 percent. However, the factor XII activity in this sample cannot be accurately quantitated due to non-parallelism. In this functional clot-based assay, factor XII activity is determined using a modified partial thromboplastin time (PTT). Non-parallelism can occur with a specific factor inhibitor directed against a clotting factor involved in the PTT reaction, inhibitor drugs such as heparin (therapy with unfractionated or low molecular weight heparin or contamination from a line), direct factor Xa inhibitors (such as rivaroxaban, apixaban, or edoxaban), direct thrombin inhibitors (such as dabigatran, argatroban, or bivalirudin), or a non-specific inhibitor such as a lupus anticoagulant. Proper interpretation requires correlation with clinical information and the results of any additional laboratory testing.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Factor XII, Activity	23-163-130737	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at: