

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 4/4/1959
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Double-Stranded DNA (dsDNA) Antibody, IgG by ELISA with Reflex to dsDNA Antibody, IgG by IFA

ARUP test code 0050215

Double-Stranded DNA (dsDNA) Ab IgG ELISA **34 IU H** (Ref Interval: 0-24)

INTERPRETIVE INFORMATION: Double-Stranded DNA (dsDNA) Ab IgG ELISA
24 IU or less.....Negative
25-30 IU.....Borderline Positive
30-60 IU.....Low Positive
60-200 IU.....Positive
201 IU or greater....Strong Positive

Positivity for anti-double stranded DNA (anti-dsDNA) IgG antibody is a diagnostic criterion of systemic lupus erythematosus (SLE). Specimens are initially screened by enzyme-linked immunosorbent assay (ELISA). If ordered as reflex (0050215), positive ELISA results (>24 IU) will be reflexed to a highly specific IFA titer (Crithidia luciliae indirect fluorescent test [CLIFT]) for confirmation. Some patients with early or inactive SLE may be positive for anti-dsDNA IgG by ELISA but negative by CLIFT. If the patient is negative by CLIFT but positive by ELISA and clinical suspicion remains, consider antinuclear antibody (ANA) testing by IFA. Additional information and recommendations for testing may be found at <https://arupconsult.com/content/systemic-lupus-erythematosus>.

Double-Stranded DNA (dsDNA) Antibody, IgG by IFA (using Crithidia luciliae)

ARUP test code 2002693

Double-Stranded DNA (dsDNA) Ab IgG IFA **1:80 *** (Ref Interval: <1:10)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

INTERPRETIVE INFORMATION: Double-Stranded DNA (dsDNA) Antibody, IgG by IFA (using Crithidia luciliae)

Positivity for anti-double stranded DNA (anti-dsDNA) IgG antibody is a diagnostic criterion of systemic lupus erythematosus (SLE). The presence of the anti-dsDNA IgG antibody is identified by IFA titer (Crithidia luciliae indirect fluorescent test [CLIFT]). CLIFT is highly specific for SLE with a sensitivity of 50-60 percent.

Some patients with early or inactive SLE may be positive for anti-dsDNA IgG by ELISA but negative by CLIFT. If the CLIFT result is negative but the patient has a positive ELISA and clinical suspicion remains, consider antinuclear antibody (ANA) testing by IFA. Additional information and recommendations for testing may be found at <http://www.arupconsult.com/Topics/AutoimmuneDz/ConnectiveTissueDz/index.html>.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Double-Stranded DNA (dsDNA) Ab IgG ELISA	23-037-109354	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Double-Stranded DNA (dsDNA) Ab IgG IFA	23-037-109354	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: