

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 11/25/1962
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Borrelia burgdorferi Antibody, IgM by Immunoblot

ARUP test code 0050253

B. burgdorferi Antibody IgM Immunoblot

Negative (Ref Interval: Negative)

Band(s) present: NONE
(Insufficient number of bands for positive result)
Current CDC recommendations for the serologic diagnosis of Lyme disease are to screen with a polyvalent EIA test and confirm equivocal and positive with Immunoblot. Both IgM and IgG Immunoblots should be performed on samples less than 4 weeks after appearance of erythema migrans. Only IgG Immunoblot should be performed on samples greater than 4 weeks after disease onset. IgM Immunoblot in the chronic stage is not recommended and does not aid in the diagnosis of neuroborreliosis or chronic Lyme disease. Please submit requests for appropriate Immunoblot testing within 10 days.

INTERPRETIVE INFORMATION: B. burgdorferi Antibody IgM Immunoblot

For this assay, a positive result is reported when any 2 or more of the following bands are present: 23, 39, or 41 kDa. All other banding patterns are reported as negative.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
B. burgdorferi Antibody IgM Immunoblot	21-090-131204	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: