

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example** 

DOB 2/26/2001 Gender: Female

**Patient Identifiers:** 01234567890ABCD, 012345

**Visit Number (FIN):** 01234567890ABCD **Collection Date:** 00/00/0000 00:00

## Herpes Simplex Virus Type 2 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050294

HSV 2 Glycoprotein G Antibody, IgG

0.04 IV

(Ref Interval: <=0.89)

REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

0.89 IV or less ...... Negative - No significant level of detectable IgG antibody to HSV

0.90 - 1.09 IV ......

type 2 glycoprotein G. Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.

1.10 IV or greater ....

Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
HSV 2 Glycoprotein G Antibody, IgG	24-022-152878	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

**END OF CHART** 

H=High, L=Low, \*=Abnormal, C=Critical

4848