

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 3/31/1968
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Scleroderma (Scl-70) (ENA) Antibody, IgG

ARUP test code 0050599

Scleroderma (Scl-70) (ENA) Antibody, IgG

2 AU/mL (Ref Interval: 0-40)
INTERPRETIVE INFORMATION: Scleroderma (Scl-70) (ENA) Ab, IgG

29 AU/mL or Less Negative
30 - 40 AU/mL Equivocal
41 AU/mL or Greater Positive

The presence of Scl-70 antibodies (also referred to as topoisomerase I, topo-I or ATA) is considered diagnostic for systemic sclerosis (SSc). Scl-70 antibodies alone are detected in about 20 percent of SSc patients and are associated with the diffuse form of the disease, which may include specific organ involvement and poor prognosis. Scl-70 antibodies have also been reported in a varying percentage of patients with systemic lupus erythematosus (SLE). Scl-70 (topo-1) is a DNA binding protein and anti-DNA/DNA complexes in the sera of SLE patients may bind to topo-I, leading to a false-positive result. The presence of Scl-70 antibody in sera may also be due to contamination of recombinant Scl-70 with DNA derived from cellular material used in immunoassays. Strong clinical correlation is recommended if both Scl-70 and dsDNA antibodies are detected.

Negative results do not necessarily rule out the presence of SSc. If clinical suspicion remains, consider further testing for centromere, RNA polymerase III and U3-RNP, PM/Scl, or Th/To antibodies.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Scleroderma (Scl-70) (ENA) Antibody, IgG	23-340-122433	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: