

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB 8/31/1993 Gender: Female

Patient Identifiers: 01234567890ABCD, 012345

Visit Number (FIN): 01234567890ABCD **Collection Date:** 00/00/0000 00:00

Cytomegalovirus Antibodies, IgG and IgM

ARUP test code 0050622

CMV Antibody IgG

0.24 U/mL (Ref Interval: <=0.70)

INTERPRETIVE INFORMATION: Cytomegalovirus Antibody, IgG

0.59 U/mL or less...... Not Detected 0.6 - 0.69 U/mL..... Indeterminate-Repeat testing in 10-14 days may be helpful.

0.70 U/mL or greater..... Detected

In immunocompromised patients, CMV serology (IgG or IgM antibody titers) may not be reliable and may be misleading in the diagnosis of acute or reactivation CMV disease. The preferred method for diagnosis is culture of virus and/or demonstration of viral antigen in peripheral white cells (buffy coat), bronchoalveolar lavage (BAL) cells, or tissue biopsies.

This test should not be used for blood donor screening, associated re-entry protocols, or for screening Human Cell, Tissues and Cellular and Tissue-Based Products (HCT/P).

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

CMV Antibody IgM

<8.0 AU/mL (Ref Interval: <=29.9)

INTERPRETIVE INFORMATION: Cytomegalovirus Antibody, IgM

29.9 AU/mL or Less Not Detected

30.0-34.9 AU/mL...... Indeterminate-Repeat testing in 10-14 days may be helpful.
35.0 AU/mL or Greater Detected-IgM antibody to CMV

detected which may indicate a current or recent infection. However, low levels of IgM antibodies may occasionally persist for more than 12 months post-infection.

A negative result does not rule out primary infection, please correlate clinically. CMV serology is not useful for the evaluation of active or reactivated infection in immunocompromised patients. Molecular diagnostic tests (i.e. PCR) are preferred in these cases.

This test should not be used for blood donor screening, associated re-entry protocols, or for screening Human Cell, Tissues and Cellular and Tissue-Based Products (HCT/P).

H=High, L=Low, *=Abnormal, C=Critical



VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
CMV Antibody IgG	24-096-126105	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
CMV Antibody IgM	24-096-126105	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical