

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 4/13/1981  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Centromere Antibody, IgG**

ARUP test code 0050714

Centromere Ab, IgG

1 AU/mL (Ref Interval: 0-40)

INTERPRETIVE INFORMATION: Centromere Ab, IgG

29 AU/mL or Less ..... Negative  
30 - 40 AU/mL ..... Equivocal  
41 AU/mL or Greater ..... Positive

When detected by this multiplex bead assay, the presence of centromere antibodies is mainly associated with CREST syndrome, a variant of systemic sclerosis (SSc). These antibodies target the centromere B, a dominant antigen of the centromeric complex associated with the centromere pattern observed in antinuclear antibody (ANA) testing by IFA. Centromere antibodies may also be seen in a varying percentage of patients with other autoimmune diseases, including diffuse cutaneous SSc, Raynaud syndrome, interstitial pulmonary fibrosis, autoimmune liver disease, systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA).

A negative result indicates no detectable IgG antibodies to centromere B. If the result is negative but clinical suspicion for SSc is strong, consider testing for ANA by IFA along with other antibodies associated with SSc, including Scl-70, U3-RNP, PM/Scl, or Th/To.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Centromere Ab, IgG	23-252-110879	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at: