

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 12/24/1969
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Centromere Antibody, IgG

ARUP test code 0050714

Centromere Ab, IgG

2 AU/mL (Ref Interval: 0-40)

INTERPRETIVE INFORMATION: Centromere Ab, IgG

29 AU/mL or Less Negative
30 - 40 AU/mL Equivocal
41 AU/mL or Greater Positive

When detected by this multiplex bead assay, the presence of centromere antibodies is mainly associated with CREST syndrome, a variant of systemic sclerosis (SSc). These antibodies target the centromere B, a dominant antigen of the centromeric complex associated with the centromere pattern observed in antinuclear antibody (ANA) testing by IFA. Centromere antibodies may also be seen in a varying percentage of patients with other autoimmune diseases, including diffuse cutaneous SSc, Raynaud syndrome, interstitial pulmonary fibrosis, autoimmune liver disease, systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA).

A negative result indicates no detectable IgG antibodies to centromere B. If the result is negative but clinical suspicion for SSc is strong, consider testing for ANA by IFA along with other antibodies associated with SSc, including Scl-70, U3-RNP, PM/Scl, or Th/To.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Centromere Ab, IgG	21-253-137819	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: