

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 11/3/1955  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Cardiolipin Antibody, IgG**

ARUP test code 0050901

Cardiolipin Antibody IgG

**>150 GPL H (Ref Interval: <=14)**

INTERPRETIVE INFORMATION: Anti-Cardiolipin IgG Ab

<=14 GPL: Negative  
15-19 GPL: Indeterminate  
20-80 GPL: Low to Moderately Positive  
81 GPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Cardiolipin Antibody IgG	23-086-109893	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

**H=High, L=Low, \*=Abnormal, C=Critical**