ARUP LABORATORIES
800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Chief Medical Officer

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB	7/5/1954
Gender:	Female
Patient Identifiers:	01234567890ABCD, 012345
Visit Number (FIN):	01234567890ABCD
Collection Date:	00/00/0000 00:00

Phosphatidylserine Antibodies, IgG, IgM, and IgA

ARUP test code 0050905

Phosphatidylserine Antibody IgG

72 GPS H (Ref Interval: 0-15)

INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgG

IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.

Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.

If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.

Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

Phosphatidylserine Antibody IgM

42 MPS H (Ref Interval: 0-21)

INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgM

The presence of IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.

Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.

If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.

Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

Phosphatidylserine Antibody IgA

24 APS H (Ref Interval: 0-19)

H=High, L=Low, *=Abnormal, C=Critical

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Phosphatidylserine Antibody IgG	24-183-123986	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine Antibody IgM	24-183-123986	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine Antibody IgA	24-183-123986	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical