

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 4/4/1981  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Phosphatidylserine Antibodies, IgG, IgM, and IgA**

ARUP test code 0050905

Phosphatidylserine Antibody IgG

**93 GPS H (Ref Interval: 0-15)**

INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgG

IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.

Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.

If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.

Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

Phosphatidylserine Antibody IgM

**64 MPS H (Ref Interval: 0-21)**

INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgM

The presence of IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.

Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.

If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.

Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

Phosphatidylserine Antibody IgA

**39 APS H (Ref Interval: 0-19)**

**H=High, L=Low, \*=Abnormal, C=Critical**

VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Phosphatidylserine Antibody IgG	22-206-401904	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine Antibody IgM	22-206-401904	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine Antibody IgA	22-206-401904	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

**H=High, L=Low, \*=Abnormal, C=Critical**

*Unless otherwise indicated, testing performed at:*

ARUP LABORATORIES | 800-522-2787 | aruplab.com  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 22-206-401904  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 2 of 2 | Printed: 9/21/2022 10:30:04 AM  
4848