

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 2/6/2006
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Humoral Immunity Panel I

ARUP test code 0050980

Diphtheria Antibody, IgG

3.5 IU/mL

INTERPRETIVE INFORMATION: Diphtheria Ab, IgG

Antibody concentration of greater than 0.1 IU/mL is usually considered protective.

Responder status is determined according to the ratio of a one month post-vaccination sample to pre-vaccination concentrations of Diphtheria IgG Abs as follows:

1. If the one month post-vaccination concentration is less than 1.0 IU/mL, the patient is considered to be a non-responder.
2. If the post-vaccination concentration is greater than or equal to 1.0 IU/mL, a patient with a ratio of less than 1.5 is a non-responder, a ratio of 1.5 to less than 3.0, a weak responder, and a ratio of 3.0 or greater, a good responder.
3. If the pre-vaccination concentration is greater than 1.0 IU/mL, it may be difficult to assess the response based on a ratio alone. A post-vaccination concentration above 2.5 IU/mL in this case is usually adequate.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Tetanus Antibody, IgG

6.6 IU/mL

H=High, L=Low, *=Abnormal, C=Critical

INTERPRETIVE INFORMATION: Tetanus Ab, IgG

Antibody concentration of greater than 0.1 IU/mL is usually considered protective.

Responder status is determined according to the ratio of a one-month post-vaccination sample to pre-vaccination concentration of Tetanus IgG Abs as follows:

1. If the one month post-vaccination concentration is less than 1.0 IU/mL, the patient is considered a non-responder.
2. If the post-vaccination concentration is greater than or equal to 1.0 IU/mL, a patient with a ratio of less than 1.5 is a non-responder, a ratio of 1.5 to less than 3.0, a weak responder, and a ratio of 3.0 or greater, a good responder.
3. If the pre-vaccination concentration is greater than 1.0 IU/mL, it may be difficult to assess the response based on a ratio alone. A post-vaccination concentration above 2.5 IU/mL in this case is usually adequate.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Pn serotype 1 IgG (P13,P20,PNX,V15) 1.19 ug/mL

Pn serotype 3 IgG (P13,P20,PNX,V15) 0.77 ug/mL

Pn serotype 4 IgG (P7,P13,P20,PNX,V15) 1.00 ug/mL

Pn serotype 5 IgG (P13,P20,PNX,V15) 9.13 ug/mL

Pn serotype 6B IgG (P7,P13,P20,PNX,V15) 9.78 ug/mL

Pn serotype 7F IgG (P13,P20,PNX,V15) 0.63 ug/mL

Pn serotype 8 IgG (P20,PNX) 0.19 ug/mL

Pn serotype 9N IgG (PNX) 0.48 ug/mL

Pn serotype 9V IgG (P7,P13,P20,PNX,V15) 0.84 ug/mL

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 24-137-153809
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 2 of 6 | Printed: 5/23/2024 10:46:10 AM
4848

Pn serotype 12F IgG (P20,PNX)	1.42 ug/mL
Pn serotype 14 IgG (P7,P13,P20,PNX,V15)	1.84 ug/mL
Pn serotype 18C IgG (P7,P13,P20,PNX,V15)	1.18 ug/mL
Pn serotype 19F IgG (P7,P13,P20,PNX,V15)	8.20 ug/mL
Pn serotype 23F IgG (P7,P13,P20,PNX,V15)	3.66 ug/mL
Pn Serotype Interpretation	See Note

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 24-137-153809
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 3 of 6 | Printed: 5/23/2024 10:46:10 AM
4848

INTERPRETIVE INFORMATION: Streptococcus pneumoniae Antibodies, IgG

A pre- and postvaccination comparison is required to adequately assess the humoral immune response to the pure polysaccharide Pneumovax 23 (PNX) and/or the protein conjugated Prevnar 7 (P7), Prevnar 13 (P13), Prevnar 20 (P20), and Vaxneuvance (V15) Streptococcus pneumoniae vaccines. Prevacination samples should be collected prior to vaccine administration. Postvaccination samples should be obtained at least 4 weeks after immunization. Testing of postvaccination samples alone will provide only general immune status of the individual to various pneumococcal serotypes.

In the case of pure polysaccharide vaccine, indication of immune system competence is further delineated as an adequate response to at least 50 percent of the serotypes in the vaccine challenge for those 2-5 years of age and to at least 70 percent of the serotypes in the vaccine challenge for those 6-65 years of age. Individual immune response may vary based on age, past exposure, immunocompetence, and pneumococcal serotype.

Responder Status	Antibody Ratio
Nonresponder	Less than twofold increase and postvaccination concentration less than 1.3 ug/mL
Good responder	At least a twofold increase and/or a postvaccination concentration greater than or equal to 1.3 ug/mL

A response to 50-70 percent or more of the serotypes in the vaccine challenge is considered a normal humoral response. (Daly, 2014) Antibody concentration greater than 1.0-1.3 ug/mL is generally considered long-term protection. (Daly, 2015)

References:

1. Daly TM, Pickering JW, Zhang X, et al. Multilaboratory assessment of threshold versus fold-change algorithms for minimizing analytical variability in multiplexed pneumococcal IgG measurements. Clin Vaccine Immunol. 2014;21(7):982-988.
2. Daly TM, Hill HR. Use and clinical interpretation of pneumococcal antibody measurements in the evaluation of humoral immune function. Clin Vaccine Immunol. 2015;22(2):148-152.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

Immunoglobulin G	1403 mg/dL	(Ref Interval: 479-1433)
Immunoglobulin A	104 mg/dL	(Ref Interval: 60-349)
Immunoglobulin M	223 mg/dL	(Ref Interval: 26-232)
Immunoglobulin G Subclass 1	720 mg/dL	(Ref Interval: 325-894)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

REFERENCE INTERVAL: Immunoglobulin G Subclass 1

The total IgG (mg/dL) can be derived from the sum of the subclass IgG1, IgG2, IgG3, and IgG4 values. However, a confirmatory and more precise total IgG is available by the turbidimetric method of quantitation for total IgG. Refer to test Immunoglobulin G, Serum (0050350).

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

Immunoglobulin G Subclass 2

326 mg/dL (Ref Interval: 156-625)

REFERENCE INTERVAL: Immunoglobulin G Subclass 2

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

Immunoglobulin G Subclass 3

88 mg/dL (Ref Interval: 34-246)

REFERENCE INTERVAL: Immunoglobulin G Subclass 3

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

Immunoglobulin G Subclass 4

189 mg/dL H (Ref Interval: 2-170)

REFERENCE INTERVAL: Immunoglobulin G Subclass 4

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

H=High, L=Low, *=Abnormal, C=Critical

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Diphtheria Antibody, IgG	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Tetanus Antibody, IgG	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 1 IgG (P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 3 IgG (P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 4 IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 5 IgG (P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 6B IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 7F IgG (P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 8 IgG (P20,PNX)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 9N IgG (PNX)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 9V IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 12F IgG (P20,PNX)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 14 IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 18C IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 19F IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn serotype 23F IgG (P7,P13,P20,PNX,V15)	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Pn Serotype Interpretation	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin G	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin A	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin M	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin G Subclass 1	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin G Subclass 2	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin G Subclass 3	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin G Subclass 4	24-137-153809	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 24-137-153809
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 6 of 6 | Printed: 5/23/2024 10:46:10 AM
4848