

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB 11/21/2019
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 01/01/2017 12:34

Humoral Immunity Panel II

ARUP test code 0050981

Diphtheria Antibody, IgG

0.0 IU/mL

INTERPRETIVE INFORMATION: Diphtheria Ab, IgG

Antibody concentration of greater than 0.1 IU/mL is usually considered protective.

Responder status is determined according to the ratio of a one month post-vaccination sample to pre-vaccination concentrations of Diphtheria IgG Abs as follows:

1. If the one month post-vaccination concentration is less than 1.0 IU/mL, the patient is considered to be a non-responder.
2. If the post-vaccination concentration is greater than or equal to 1.0 IU/mL, a patient with a ratio of less than 1.5 is a non-responder, a ratio of 1.5 to less than 3.0, a weak responder, and a ratio of 3.0 or greater, a good responder.
3. If the pre-vaccination concentration is greater than 1.0 IU/mL, it may be difficult to assess the response based on a ratio alone. A post-vaccination concentration above 2.5 IU/mL in this case is usually adequate.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

Tetanus Antibody, IgG

0.0 IU/mL

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

INTERPRETIVE INFORMATION: Tetanus Ab, IgG

Antibody concentration of greater than 0.1 IU/mL is usually considered protective.

Responder status is determined according to the ratio of a one-month post-vaccination sample to pre-vaccination concentration of Tetanus IgG Abs as follows:

1. If the one month post-vaccination concentration is less than 1.0 IU/mL, the patient is considered a non-responder.
2. If the post-vaccination concentration is greater than or equal to 1.0 IU/mL, a patient with a ratio of less than 1.5 is a non-responder, a ratio of 1.5 to less than 3.0, a weak responder, and a ratio of 3.0 or greater, a good responder.
3. If the pre-vaccination concentration is greater than 1.0 IU/mL, it may be difficult to assess the response based on a ratio alone. A post-vaccination concentration above 2.5 IU/mL in this case is usually adequate.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

Pneumo serotype 1 IgG (P13,PNX)	0.07 ug/mL
Pneumo serotype 3 IgG (P13,PNX)	0.45 ug/mL
Pneumo serotype 4 IgG (P7,P13,PNX)	0.04 ug/mL
Pneumo serotype 5 IgG (P13,PNX)	0.60 ug/mL
Pneumo serotype 6B IgG (P7,P13,PNX)	0.20 ug/mL
Pneumo serotype 7F IgG (P13,PNX)	1.40 ug/mL
Pneumo serotype 8 IgG (PNX)	0.09 ug/mL
Pneumo serotype 9N IgG (PNX)	0.08 ug/mL
Pneumo serotype 9V IgG (P7,P13,PNX)	0.03 ug/mL

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-153-102165
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 2 of 4 | Printed: 12/7/2020 11:24:54 AM
4848

Pneumo serotype 12F IgG (PNX) 0.04 ug/mL

Pneumo serotype 14 IgG (P7,P13,PNX) 0.18 ug/mL

Pneumo serotype 18C IgG (P7,P13,PNX) 0.13 ug/mL

Pneumo serotype 19F IgG (P7,P13,PNX) 0.37 ug/mL

Pneumo serotype 23F IgG (P7,P13,PNX) 0.04 ug/mL

Pneumo Serotype Interpretation

See Note

INTERPRETIVE INFORMATION: Streptococcus pneumoniae Antibodies, IgG

A pre- and post-vaccination comparison is required to adequately assess the humoral immune response to Prevnar 7 (P7), Prevnar 13 (P13), and/or Pneumovax 23 (PNX) Streptococcus pneumoniae vaccines. Pre-vaccination samples should be collected prior to vaccine administration. Post-vaccination samples should be obtained at least 4 weeks after immunization. Testing of post-vaccination samples alone will provide only general immune status of the individual to various pneumococcal serotypes.

In the case of pure polysaccharide vaccine, indication of immune system competence is further delineated as an adequate response to at least 50 percent of the serotypes in the vaccine challenge for those 2-5 years of age and to at least 70 percent of the serotypes in the vaccine challenge for those 6-65 years of age. Individual immune response may vary based on age, past exposure, immunocompetence, and pneumococcal serotype.

Responder Status	Antibody Ratio
Non-Responder	Less than 2-fold
Weak Responder	2-fold to 4-fold
Good Responder	Greater than 4-fold

A response to 50-70 percent or more of the serotypes in the vaccine challenge is considered a normal humoral response(1). Antibody concentration greater than 1.0 - 1.3 ug/mL is generally considered long-term protection(2).

- References:
- Daly TM, Pickering JW, Zhang X, Prince HE, Hill HR. Multilaboratory assessment of threshold versus fold-change algorithms for minimizing analytical variability in multiplexed pneumococcal IgG measurements. Clin Vaccine Immunol. 2014;21(7):982-8.
 - Daly TM, Hill HR. Use and Clinical Interpretation of Pneumococcal Antibody Measurements in the Evaluation of Humoral Immune Function. Clin Vaccine Immunol. 2015;22(2):148-152.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-153-102165
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 3 of 4 | Printed: 12/7/2020 11:24:54 AM
4848

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Diphtheria Antibody, IgG	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/4/2020 11:19:00 AM
Tetanus Antibody, IgG	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/4/2020 11:19:00 AM
Pneumo serotype 1 IgG (P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 3 IgG (P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 4 IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 5 IgG (P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 6B IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 7F IgG (P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 8 IgG (PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 9N IgG (PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 9V IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 12F IgG (PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 14 IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 18C IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 19F IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo serotype 23F IgG (P7,P13,PNX)	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM
Pneumo Serotype Interpretation	20-153-102165	6/1/2020 5:50:00 AM	6/2/2020 12:04:39 AM	6/3/2020 5:27:00 AM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-153-102165
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 4 of 4 | Printed: 12/7/2020 11:24:54 AM
4848