

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 10/17/1989
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 01/01/2017 12:34

Cardiolipin Antibodies, IgG, IgM, and IgA

ARUP test code 0051162

Cardiolipin Antibody IgG

3 GPL (Ref Interval: 0-14)

INTERPRETIVE INFORMATION: Anti-Cardiolipin IgG Ab

0-14 GPL: Negative
15-19 GPL: Indeterminate
20-80 GPL: Low to Moderately Positive
81 GPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units or greater than 99th percentile) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

Cardiolipin Antibody IgM

9 MPL (Ref Interval: 0-12)

INTERPRETIVE INFORMATION: Anti-Cardiolipin IgM

0-12 MPL: Negative
13-19 MPL: Indeterminate
20-80 MPL: Low to Moderately Positive
81 MPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units or greater than 99th percentile) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 19-291-128020
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 1 of 2 | Printed: 12/7/2020 11:25:17 AM
4848

Cardiolipin Antibody IgA

2 APL

(Ref Interval: 0-11)

INTERPRETIVE INFORMATION: Cardiolipin Antibodies, IgA

0-11 APL: Negative

12-19 APL: Indeterminate

20-80 APL: Low to Moderately Positive

81 APL or above: High Positive

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Cardiolipin Antibody IgG	19-291-128020	10/18/2019 11:45:00 AM	10/20/2019 9:50:22 AM	10/20/2019 11:23:00 PM
Cardiolipin Antibody IgM	19-291-128020	10/18/2019 11:45:00 AM	10/20/2019 9:50:22 AM	10/20/2019 11:23:00 PM
Cardiolipin Antibody IgA	19-291-128020	10/18/2019 11:45:00 AM	10/20/2019 9:50:22 AM	10/20/2019 11:23:00 PM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 19-291-128020
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 2 of 2 | Printed: 12/7/2020 11:25:17 AM
4848