

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 1/8/1950  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**F-Actin (Smooth Muscle) Antibody, IgG by ELISA with Reflex to Smooth Muscle Antibody, IgG Titer**

ARUP test code 0051174

F-Actin (Smooth Muscle) Ab, IgG by ELISA

8 Units

(Ref Interval: 0-19)

If F-Actin (Smooth Muscle) Antibody, IgG is negative, the Smooth Muscle Antibody titer by IFA is not performed.

REFERENCE INTERVAL: F-Actin (Smooth Muscle) Antibody, IgG by ELISA

- 19 Units or less ..... Negative
- 20 - 30 Units ..... Weak Positive-Suggest repeat testing in two to three weeks with fresh specimen.
- 31 Units or greater..... Positive-Suggestive of autoimmune hepatitis type 1 or chronic active hepatitis.

F-actin IgG antibodies have been shown to have increased sensitivity for autoimmune hepatitis (AIH) but lower specificity than smooth muscle antibodies (SMA). F-actin IgG antibodies can also be seen in SMA-negative disease controls (non-AIH), especially in patients with primary biliary cirrhosis and chronic hepatitis C infections. Some patients with AIH may be SMA-positive but negative for F-actin IgG. Consider testing for SMA by IFA if suspicion for AIH is strong.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
F-Actin (Smooth Muscle) Ab, IgG by ELISA	21-023-102416	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: