

Patient Report | FINAL

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108

Physician: Doctor, Example

UNITED STATES

Patient: Patient, Example

DOB 10/6/1951 **Gender:** Male

Patient Identifiers: 01234567890ABCD, 012345

Visit Number (FIN): 01234567890ABCD **Collection Date:** 00/00/0000 00:00

Motor Neuropathy Panel

ARUP test code 0051225

MAG Antibody, IgM Elisa

<1000 TU

(Ref Interval: 0-999)

H=High, L=Low, *=Abnormal, C=Critical

4848



	INTERPRETIVE INFORMATION: MAG Antibody, IgM ELISA			
	An elevated IgM antibody concentration greater than 999 TU against myelin-associated glycoprotein (MAG) suggests active demyelination in peripheral neuropathy. A normal concentration (less than 999 TU) generally rules out an anti-MAG antibody-associated peripheral neuropathy. TU=Titer Units This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.			
SGPG Antibody, IgM	0.09 IV	(Ref Interval: 0.00-0.99)		
	INTERPRETIVE INFORMATION: SGPG Antibody, IgM			
	The majority of sulfate-3-glucuronyl paragloboside (SGPG) IgM-positive sera will show reactivity against MAG. Patients who are SGPG IgM positive and MAG IgM negative may have multi-focal motor neuropathy with conduction block. This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.			
Asialo-GM1 Antibodies, IgG/IgM	8 IV	(Ref Interval: 0-50)		
GM1 Antibodies, IgG/IgM	26 IV	(Ref Interval: 0-50)		
GD1a Antibodies, IgG/IgM	8 IV	(Ref Interval: 0-50)		
GD1b Antibodies, IgG/IgM	8 IV	(Ref Interval: 0-50)		
GQ1b Antibodies, IgG/IgM	8 IV	(Ref Interval: 0-50)		



INTERPRETIVE INFORMATION: Ganglioside (Asialo-GM1, GM1, GM2, GD1a, GD1b, and GQ1b) Antibodies, IgG/IgM

29 IV or less: Negative 30-50 IV: Equivocal 51-100 IV: Positive

101 IV or greater: Strong Positive

Ganglioside antibodies are associated with diverse peripheral neuropathies. Elevated antibody levels to ganglioside-monosialic acid (GM1), and the neutral glycolipid, asialo GM1 are associated with motor or sensorimotor neuropathies, particularly multifocal motor neuropathy. Anti-GM1 may occur as IgM (polyclonal or monoclonal) or IgG antibodies. These antibodies may also be found in patients with diverse connective tissue diseases as well as normal individuals. GD1a antibodies are associated with different variants of Guillain-Barre syndrome (GBS) particularly acute motor axonal neuropathy while GD1b antibodies are predominantly found in sensory ataxic neuropathy syndrome. Anti-GQ1b antibodies are seen in more than 80 percent of patients with Miller-Fisher syndrome and may be elevated in GBS patients with ophthalmoplegia. The role of isolated anti-GM2 antibodies is unknown. These tests by themselves are not diagnostic and should be used in conjunction with other clinical parameters to confirm disease.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Immunoglobulin G	1222 mg/dL	(Ref Interval: 768-1632)
Immunoglobulin A	137 mg/dL	(Ref Interval: 68-408)
Immunoglobulin M	75 mg/dL	(Ref Interval: 35-263)
Total Protein, Serum	7.0 g/dL	(Ref Interval: 6.3-8.2)
Albumin	4.55 g/dL	(Ref Interval: 3.75-5.01)
Alpha 1 Globulin	0.22 g/dL	(Ref Interval: 0.19-0.46)
Alpha 2 Globulin	0.48 g/dL	(Ref Interval: 0.48-1.05)
Beta Globulin	0.66 g/dL	(Ref Interval: 0.48-1.10)
Gamma	1.09 g/dL	(Ref Interval: 0.62-1.51)



Monoclonal Protein	Not Applicable g/dL		
Immunofixation	IFE Done		
SPEP/IFE Interpretation	See Note Normal SPEP pattern. IFE gel shows a normal pattern; no monoclonal proteins seen.		
EER Motor Neuropathy Panel	See Note Authorized individuals can access the ARUP Enhanced Report using the following link:		



VERIFIED/REPORTED DATES							
Procedure	Accession	Collected	Received	Verified/Reported			
MAG Antibody, IgM Elisa	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
SGPG Antibody, IgM	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Asialo-GM1 Antibodies, IgG/IgM	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
GM1 Antibodies, IgG/IgM	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
GD1a Antibodies, IgG/IgM	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
GD1b Antibodies, IgG/IgM	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
GQ1b Antibodies, IgG/IgM	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Immunoglobulin G	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Immunoglobulin A	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Immunoglobulin M	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Total Protein, Serum	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Albumin	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Alpha 1 Globulin	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Alpha 2 Globulin	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Beta Globulin	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Gamma	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Monoclonal Protein	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
Immunofixation	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
SPEP/IFE Interpretation	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			
EER Motor Neuropathy Panel	24-043-129373	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00			

END OF CHART