

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 9/25/1978  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG with Reflex to Type 1 and 2 Glycoprotein G-Specific Ab, IgG**

ARUP test code 0051708

HSV Type 1/2 Combined Ab, IgG >22.40 IV

Specimen tested positive for Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG. ARUP test codes 0050292 and 0050294 will be added. Additional charges apply.

INTERPRETIVE INFORMATION: HSV 1/2 COMBINED Ab SCREEN, IgG  
0.89 IV or less.....Not Detected  
0.90-1.09 IV.....Indeterminate- Repeat testing in 10-14 days may be helpful.  
1.10 IV or greater.....Detected

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

**Herpes Simplex Virus Type 1 Glycoprotein G-Specific Antibody, IgG by CIA**

ARUP test code 0050292

HSV 1 Glycoprotein G Ab, IgG **56.00 IV H** (Ref Interval: <=0.89)

REFERENCE INTERVAL: HSV 1 Glycoprotein G Ab, IgG

0.89 IV or less ..... Negative - No significant level of detectable IgG antibody to HSV type 1 glycoprotein G.  
0.90 - 1.09 IV ..... Equivocal - Questionable presence of IgG antibody to HSV type 1 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.  
1.10 IV or greater ... Positive - IgG antibody to HSV type 1 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

**Herpes Simplex Virus Type 2 Glycoprotein G-Specific Antibody, IgG by CIA**

ARUP test code 0050294

**H=High, L=Low, \*=Abnormal, C=Critical**

HSV 2 Glycoprotein G Antibody, IgG

**1.52 IV H (Ref Interval: <=0.89)**

REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

- 0.89 IV or less ..... Negative - No significant level of detectable IgG antibody to HSV type 2 glycoprotein G.
- 0.90 - 1.09 IV ..... Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
- 1.10 IV or greater .... Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
HSV Type 1/2 Combined Ab, IgG	22-322-402198	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
HSV 1 Glycoprotein G Ab, IgG	22-322-402198	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
HSV 2 Glycoprotein G Antibody, IgG	22-322-402198	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: