

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 5/25/1995
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG with Reflex to Type 1 and 2 Glycoprotein G-Specific Ab, IgG

ARUP test code 0051708

HSV Type 1/2 Combined Ab, IgG >22.40 IV

Specimen tested positive for Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG. ARUP test codes 0050292 and 0050294 will be added. Additional charges apply.

INTERPRETIVE INFORMATION: HSV 1/2 COMBINED Ab SCREEN, IgG
0.89 IV or less.....Not Detected
0.90-1.09 IV.....Indeterminate- Repeat testing
in 10-14 days may be helpful.
1.10 IV or greater.....Detected

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Herpes Simplex Virus Type 1 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050292

HSV 1 Glycoprotein G Ab, IgG 0.04 IV (Ref Interval: <=0.90)

INTERPRETIVE INFORMATION: HSV 1 Glycoprotein G Ab, IgG (CIA)

0.90 IV or less Negative - No significant level of detectable IgG antibody to HSV type 1 glycoprotein G.
0.91 - 1.09 IV Equivocal - Questionable presence of IgG antibody to HSV type 1 glycoprotein G. Repeat testing in 10-14 days may be helpful.
1.10 IV or greater Positive - IgG antibody to HSV type 1 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type specific HSV antigens 1 and 2 in the early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

Herpes Simplex Virus Type 2 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050294

H=High, L=Low, *=Abnormal, C=Critical



HSV 2 Glycoprotein G Antibody, IgG

12.10 IV H (Ref Interval: <=0.90)

INTERPRETIVE INFORMATION: HSV 2 Glycoprotein G Ab, IgG (CIA)

- 0.90 IV or less Negative - No significant level of detectable IgG antibody to HSV type 2 glycoprotein G.
- 0.91 - 1.09 IV Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10-14 days may be helpful.
- 1.10 IV or greater Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type specific HSV antigens 1 and 2 in the early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
HSV Type 1/2 Combined Ab, IgG	18-247-400780	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
HSV 1 Glycoprotein G Ab, IgG	18-247-400780	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
HSV 2 Glycoprotein G Antibody, IgG	18-247-400780	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical