

Client: Example Client ABC123  
 123 Test Drive  
 Salt Lake City, UT 84108  
 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 01/01/1990  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 01/01/2017 12:34

**Treponema pallidum Antibody, IgG by IFA (CSF)**

ARUP test code 0055273

Fluorescent Treponema Antibody CSF

Minimal React

(Ref Interval: Non Reactive)

In the absence of historical or clinical evidence of treponemal infection, a minimal reactive FTA result should be considered equivocal. A second specimen should be obtained 1-2 weeks after the initial specimen and submitted to the laboratory for serologic testing.

**INTERPRETIVE INFORMATION:** Treponema pallidum Antibody, IgG by IFA (CSF)

The significance of a reactive result in the FTA-ABS CSF test is unknown. The CSF from persons treated in the secondary or latent stage of syphilis and without signs of neurosyphilis may be reactive. A nonreactive result in the FTA-ABS CSF test suggests the absence of neurosyphilis.

Treponema pallidum (VDRL), Cerebrospinal Fluid with Reflex to Titer (0050206) is the recommended test for CSF specimens. If suspicion of neurosyphilis remains after VDRL testing, testing of the CSF with FTA-ABS may be considered.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Fluorescent Treponema Antibody CSF	15-237-402887	8/22/2015 2:55:00 AM	8/26/2015 6:13:40 PM	8/29/2015 11:43:24 AM

END OF CHART

**H - high L - low \* - abnormal C - critical**