

Client: Example Client ABC123  
 123 Test Drive  
 Salt Lake City, UT 84108  
 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 5/16/2004  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Tissue Transglutaminase Antibody, IgG**

ARUP test code 0056009

Tissue Transglutaminase Antibody,IgG

1 U/mL (Ref Interval: 0-5)  
 INTERPRETIVE INFORMATION: Tissue Transglutaminase Ab, IgG

5 U/mL or less: ..... Negative  
 6-9 U/mL: ..... Weak Positive  
 10 U/mL or greater: ..... Positive

The tTG IgG assay may aid in the diagnosis of gluten-sensitivity enteropathy (i.e., celiac disease, dermatitis herpetiformis) in tTG IgA negative patients with confirmed IgA deficiency. A negative tTG IgG test alone does not rule out gluten-sensitive enteropathy.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Tissue Transglutaminase Antibody,IgG	19-228-103485	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical