

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB 2/22/1987
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Wound Culture and Gram Stain

ARUP test code 0060132

Collected: 00/00/0000 00:00 MT
Started: 00/00/0000 00:00 MT

Source: **wound**

Body Site: **Toe**

Free Text Sources:

Stains/Preparations

Gram Stain

1+ Gram Positive Cocci
No polymorphonuclear leukocytes (PMNs) seen

Final Report

2+ Staphylococcus aureus
2+ Coagulase negative Staphylococcus species

Susceptibility Results

Organism: Staphylococcus aureus

Clindamycin	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=0.5
Daptomycin	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=1

S=Susceptible, I=Intermediate, R=Resistant, NonS=Nonsusceptible, IND=Indeterminate, SDD=Susceptibility is dose dependent, None=Interpretive guidelines are not available

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 23-173-124751
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 1 of 2 | Printed: 6/26/2023 9:41:57 AM
4848

Erythromycin	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=0.5
Gentamicin	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=2
Levofloxacin	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=0.5
Linezolid	Interpretation: SUSCEPTIBLE MIC (ug/mL): 1
Oxacillin	Interpretation: RESISTANT MIC (ug/mL): >=4
Penicillin	Interpretation: RESISTANT MIC (ug/mL): >=2
Rifampin	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=0.5
Tetracycline	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=0.5
Trimethoprim/Sulfamethoxazole	Interpretation: SUSCEPTIBLE MIC (ug/mL): <=1/19
Vancomycin	Interpretation: SUSCEPTIBLE MIC (ug/mL): 1

Interpretive Information Interpretation: **SEE NOTE**

Gentamicin: susceptible- For staphylococci that test susceptible, gentamicin is used only in combination with other active agents that test susceptible.

Rifampin should not be used alone for antimicrobial therapy.

VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Wound Culture and Gram Stain	23-173-124751	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

S=Susceptible, I=Intermediate, R=Resistant, NonS=Nonsusceptible, IND=Indeterminate, SDD=Susceptibility is dose dependent, None=Interpretive guidelines are not available

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: