

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 9/14/1974  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Prostate Specific Antigen, Free Percentage (Includes Free PSA and Total PSA)**

ARUP test code 0080206

**Prostate Specific Antigen, Total** 0.7 ng/mL (Ref Interval: 0.0-4.0)  
INTERPRETIVE INFORMATION: Prostate Specific Antigen

The Roche PSA electrochemiluminescent immunoassay was used. Results obtained with different test methods or kits cannot be used interchangeably. The Roche PSA method is approved for use as an aid in the detection of prostate cancer when used in conjunction with a digital rectal exam in men age 50 and older. The Roche PSA method is also indicated for the serial measurement of PSA to aid in the prognosis and management of prostate cancer patients. Elevated PSA concentrations can only suggest the presence of prostate cancer until biopsy is performed. PSA concentrations can also be elevated in benign prostatic hyperplasia or inflammatory conditions of the prostate. PSA is generally not elevated in healthy men or men with non-prostatic carcinoma.

**Prostate Specific Antigen, Free** 0.1 ng/mL

**Prostate Specific Antigen, Percent Free** 14 %  
INTERPRETIVE INFORMATION: Prostate Specific Antigen, Free Percentage

ARUP uses the Roche Free PSA electrochemiluminescent immunoassay method in conjunction with the Roche PSA electrochemiluminescent immunoassay method to determine the free PSA percentage. Values obtained with different assay methods should not be used interchangeably. The free PSA percentage is an aid in distinguishing prostate cancer from benign prostatic conditions in men age 50 and older with a total PSA between 3 and 10 ng/mL and negative digital rectal examination findings. Prostatic biopsy is required for the diagnosis of cancer.

In patients with total PSA concentrations of 4-10 ng/mL, the probability of finding prostate cancer on needle biopsy by age in years is:

%fPSA	50-59	60-69	70 or older
0 - 10%	49%	58%	65%
11 - 18%	27%	34%	41%
19 - 25%	18%	24%	30%
Greater than 25%	9%	12%	16%

Other factors may help determine the actual risk of prostate cancer in individual patients.

**H=High, L=Low, \*=Abnormal, C=Critical**

VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Prostate Specific Antigen, Total	20-361-108008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Prostate Specific Antigen, Free	20-361-108008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Prostate Specific Antigen, Percent Free	20-361-108008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 20-361-108008  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 2 of 2 | Printed: 12/28/2020 11:40:31 AM  
4848