

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 5/21/1952  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 01/01/2017 12:34

**Cardiolipin Antibodies, IgG and IgM**

ARUP test code 0099344

**Cardiolipin Antibody IgG**

**67 GPL H (Ref Interval: 0-14)**

INTERPRETIVE INFORMATION: Anti-Cardiolipin IgG Ab

0-14 GPL: Negative  
15-19 GPL: Indeterminate  
20-80 GPL: Low to Moderately Positive  
81 GPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units or greater than 99th percentile) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

**Cardiolipin Antibody IgM**

**>150 MPL H (Ref Interval: 0-12)**

INTERPRETIVE INFORMATION: Anti-Cardiolipin IgM

0-12 MPL: Negative  
13-19 MPL: Indeterminate  
20-80 MPL: Low to Moderately Positive  
81 MPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units or greater than 99th percentile) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at:

**ARUP LABORATORIES | 800-522-2787 | aruplab.com**  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 19-130-117978  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 1 of 2 | Printed: 12/7/2020 11:28:35 AM  
4848

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Cardiolipin Antibody IgG	19-130-117978	5/10/2019 11:41:00 AM	5/11/2019 1:46:30 AM	5/11/2019 4:28:00 PM
Cardiolipin Antibody IgM	19-130-117978	5/10/2019 11:41:00 AM	5/11/2019 1:46:30 AM	5/11/2019 4:28:00 PM

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

**ARUP LABORATORIES | 800-522-2787 | aruplab.com**  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 19-130-117978  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 2 of 2 | Printed: 12/7/2020 11:28:35 AM  
4848