

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 6/15/1966
Gender: Unknown
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

RNA Polymerase III Antibody, IgG

ARUP test code 2001601

RNA Polymerase III Antibody, IgG

6 Units (Ref Interval: 0-19)
INTERPRETIVE INFORMATION: RNA Polymerase III Antibody, IgG

19 Units or lessNegative
20 - 39 UnitsWeak Positive
40 - 80 UnitsModerate Positive
81 Units or greater ...Strong Positive

The presence of RNA polymerase III IgG antibody, when considered in conjunction with other laboratory and clinical findings, is an aid in the diagnosis of systemic sclerosis (SSc) with increased incidence of skin involvement and renal crisis with the diffuse cutaneous form of SSc. RNA polymerase III IgG antibody occur in about 11-23 percent of SSc patients, and typically in the absence of anti-centromere and anti-Scl-70 antibodies.

A negative result indicates no detectable IgG antibodies to the dominant antigen of RNA polymerase III and does not rule out the possibility of SSc. False-positive results may also occur due to non-specific binding of immune complexes. Strong clinical correlation is recommended.

If clinical suspicion remains, consider additional testing for other antibodies associated with SSc, including centromere, Scl-70, U3-RNP, PM/Scl, or Th/To.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
RNA Polymerase III Antibody, IgG	21-021-108841	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: