

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example** 

**DOB** 7/12/1984

**Gender:** Male

**Patient Identifiers:** 01234567890ABCD, 012345

**Visit Number (FIN):** 01234567890ABCD **Collection Date:** 00/00/0000 00:00

## Inhibitor Assay, PTT with Reflex to PTT 1:1 Mix, with Reflex to 1-Hour Incubation

ARUP test code 2003266

PTT, Inhibitor Screen

**142** sec H

(Ref Interval: 32-48)

PTT 1:1 mixing study added.

A PTT-based inhibitor is detected.
Possibilities include a specific factor inhibitor, lupus anticoagulant, anticoagulant medications such as heparin (therapy with unfractionated or low molecular weight heparin or contamination from a line), direct thrombin inhibitors, or factor Xa inhibitors, dysfibrinogenemia, or monoclonal proteins. This pattern can occasionally be seen in the presence of multiple or severe factor deficiencies. Please correlate clinically. If a lupus anticoagulant is suspected, recommend the lupus anticoagulant reflexive panel for confirmation.

## Partial Thromboplastin Time 1:1 Mix (performed if PTT > 48 seconds)

ARUP test code 2003270

PTT, Inhibitor Screen, 1:1 Mix

97 sec H

(Ref Interval: 32-48)

The PTT 1:1 mixing study is prolonged; therefore, no further testing added.

H=High, L=Low, \*=Abnormal, C=Critical

4848



VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
PTT, Inhibitor Screen	23-031-142006	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
PTT, Inhibitor Screen, 1:1 Mix	23-031-142006	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

4848