

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 11/8/1948
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 01/01/2017 12:34

Rheumatoid Arthritis Panel with Reflex to Rheumatoid Factors, IgA, IgG, and IgM by ELISA

ARUP test code 2003278

Rheumatoid Factor	56 IU/mL	H	(Ref Interval: 0-14)
Cyclic Citrullinated Peptide Ab, IgG	214 Units	H	(Ref Interval: 0-19)

INTERPRETIVE INFORMATION: Cyclic Citrullinated Peptide Antibody, IgG

19 Units or less	Negative
20-39 Units	Weak Positive
40-59 Units	Moderate Positive
60 Units or greater	Strong Positive

Anti-cyclic citrullinated peptide (anti-CCP), IgG antibodies are present in about 69-83 percent of patients with rheumatoid arthritis (RA) and have specificities of 93-95 percent. These autoantibodies may be present in the preclinical phase of disease, are associated with future RA development, and may predict radiographic joint destruction. Patients with weak positive results should be monitored and testing repeated.

Rheumatoid Factors, IgA, IgG, and IgM by ELISA

ARUP test code 0051298

Rheumatoid Factor, IgA by ELISA	29 Units	H	(Ref Interval: 0-6)
Rheumatoid Factor, IgG by ELISA	7 Units	H	(Ref Interval: 0-6)

INTERPRETIVE INFORMATION: Rheumatoid Factor, IgA by ELISA

The presence of all three rheumatoid factor (RF) isotypes at abnormal levels has high specificity for a diagnosis of rheumatoid arthritis (RA). However, the presence of RF isotypes in any combination may be found in a variety of conditions, including Sjogren syndrome and hepatitis infections.

INTERPRETIVE INFORMATION: Rheumatoid Factor, IgG by ELISA

The presence of all three rheumatoid factor (RF) isotypes at abnormal levels has high specificity for a diagnosis of rheumatoid arthritis (RA). However, the presence of RF isotypes in any combination may be found in a variety of conditions, including Sjogren syndrome and hepatitis infections.

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-175-118276
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 1 of 2 | Printed: 12/7/2020 11:30:03 AM
4848

Rheumatoid Factor, IgM by ELISA

>100 Units H (Ref Interval: 0-6)

INTERPRETIVE INFORMATION: Rheumatoid Factor, IgM by ELISA

The presence of all three rheumatoid factor (RF) isotypes at abnormal levels has high specificity for a diagnosis of rheumatoid arthritis (RA). However, the presence of RF isotypes in any combination may be found in a variety of conditions, including Sjogren syndrome and hepatitis infections.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Rheumatoid Factor	20-175-118276	6/23/2020 1:30:00 PM	6/24/2020 10:56:14 PM	6/25/2020 4:07:00 AM
Rheumatoid Factor, IgA by ELISA	20-175-118276	6/23/2020 1:30:00 PM	6/25/2020 4:06:57 AM	6/26/2020 2:17:00 PM
Rheumatoid Factor, IgG by ELISA	20-175-118276	6/23/2020 1:30:00 PM	6/25/2020 4:06:57 AM	6/26/2020 2:17:00 PM
Rheumatoid Factor, IgM by ELISA	20-175-118276	6/23/2020 1:30:00 PM	6/25/2020 4:06:57 AM	6/26/2020 2:17:00 PM
Cyclic Citrullinated Peptide Ab, IgG	20-175-118276	6/23/2020 1:30:00 PM	6/24/2020 10:56:14 PM	6/25/2020 5:10:00 PM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-175-118276
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 2 of 2 | Printed: 12/7/2020 11:30:03 AM
4848