

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 4/12/1982
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Ganglioside (GM1, GD1b, and GQ1b) Antibodies, IgG and IgM

ARUP test code 2004998

GM1 Antibody, IgG	7 IV		(Ref Interval: 0-50)
GM1 Antibody, IgM	5 IV		(Ref Interval: 0-50)
<p>INTERPRETIVE INFORMATION: Ganglioside (GM1) Antibodies, IgG and IgM</p> <p>29 IV or less: Negative 30-50 IV: Equivocal 51-100 IV: Positive 101 IV or greater: Strong positive</p> <p>Ganglioside antibodies are associated with diverse peripheral neuropathies. Elevated antibody levels to ganglioside-monosialic acid (GM1) are associated with motor or sensorimotor neuropathies, particularly multifocal motor neuropathy. Anti-GM1 may occur as IgM (polyclonal or monoclonal) or IgG antibodies. These antibodies may also be found in patients with diverse connective tissue diseases as well as normal individuals. These tests by themselves are not diagnostic and should be used in conjunction with other clinical parameters to confirm disease.</p> <p>This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.</p>			
GD1b Antibody, IgG	305 IV	H	(Ref Interval: 0-50)
GD1b Antibody, IgM	59 IV	H	(Ref Interval: 0-50)
GQ1b Antibody, IgG	312 IV	H	(Ref Interval: 0-50)
GQ1b Antibody, IgM	76 IV	H	(Ref Interval: 0-50)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

INTERPRETIVE INFORMATION: Ganglioside (GD1b and GQ1b) Antibodies, IgG and IgM

29 IV or less: Negative
30-50 IV: Equivocal
51-100 IV: Positive
101 IV or greater: Strong positive

Ganglioside antibodies are associated with diverse peripheral neuropathies. GD1b antibodies are predominantly found in sensory ataxic neuropathy syndrome. GQ1b antibodies are seen in more than 80% of patients with Miller-Fisher syndrome and may be elevated in Guillain-Barre syndrome patients with ophthalmoplegia. These tests by themselves are not diagnostic and should be used in conjunction with other clinical parameters to confirm disease.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
GM1 Antibody, IgG	24-015-400921	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GM1 Antibody, IgM	24-015-400921	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GD1b Antibody, IgG	24-015-400921	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GD1b Antibody, IgM	24-015-400921	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GQ1b Antibody, IgG	24-015-400921	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GQ1b Antibody, IgM	24-015-400921	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: