

Client: Example Client ABC123 123 Test Drive

Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example** 

**DOB** 2/5/1964 **Sex:** Male

**Patient Identifiers:** 01234567890ABCD, 012345

**Visit Number (FIN):** 01234567890ABCD **Collection Date:** 01/01/2017 12:34

## Phosphatidylserine Antibodies, IgG and IgM

ARUP test code 2006495

Phosphatidylserine Antibody IgG

43 U/mL H

(Ref Interval: 0-10)

INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgG

The presence of IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.

Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.

If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.

Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

Phosphatidylserine Antibody IgM

34 U/mL H

(Ref Interval: 0-24)

INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgM

The presence of IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.

Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.

If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.

Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Phosphatidylserine Antibody IgG	20-162-112080	6/10/2020 9:25:00 AM	6/11/2020 10:30:29 AM	6/13/2020 3:02:00 PM
Phosphatidylserine Antibody IgM	20-162-112080	6/10/2020 9:25:00 AM	6/11/2020 10:30:29 AM	6/15/2020 4:43:00 PM

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical