

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 9/22/1999
Sex: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Motor and Sensory Neuropathy Evaluation with Immunofixation Electrophoresis and Reflex to Titer and Neuronal Immunoblot

ARUP test code 2007967

MAG Antibody, IgM Elisa

<1000 TU (Ref Interval: 0-999)

INTERPRETIVE INFORMATION: MAG Antibody, IgM ELISA

An elevated IgM antibody concentration greater than 999 TU against myelin-associated glycoprotein (MAG) suggests active demyelination in peripheral neuropathy. A normal concentration (less than 999 TU) generally rules out an anti-MAG antibody-associated peripheral neuropathy.

TU=Titer Units

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

SGPG Antibody, IgM

0.04 IV (Ref Interval: 0.00-0.99)

INTERPRETIVE INFORMATION: SGPG Antibody, IgM

The majority of sulfate-3-glucuronyl paragloboside (SGPG) IgM-positive sera will show reactivity against MAG. Patients who are SGPG IgM positive and MAG IgM negative may have multi-focal motor neuropathy with conduction block.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Purkinje Cell/Neuronal Nuclear IgG Scrn

None Detected (Ref Interval: None Detected)

ANNA-1, ANNA-2, PCCA-1 or PCCA-Tr(DNER) antibodies not detected. No further testing will be performed.

INTERPRETIVE INFORMATION: Purkinje Cell/Neuronal Nuclear IgG Scrn

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

performed in a CLIA certified laboratory and is intended for clinical purposes.

Asialo-GM1 Antibodies, IgG/IgM	11 IV	(Ref Interval: 0-50)
GM1 Antibodies, IgG/IgM	8 IV	(Ref Interval: 0-50)
GD1a Antibodies, IgG/IgM	5 IV	(Ref Interval: 0-50)
GD1b Antibodies, IgG/IgM	5 IV	(Ref Interval: 0-50)
GQ1b Antibodies, IgG/IgM	5 IV	(Ref Interval: 0-50)

INTERPRETIVE INFORMATION: Ganglioside (Asialo-GM1, GM1, GM2, GD1a, GD1b, and GQ1b) Antibodies, IgG/IgM

29 IV or less: Negative
30-50 IV: Equivocal
51-100 IV: Positive
101 IV or greater: Strong Positive

Ganglioside antibodies are associated with diverse peripheral neuropathies. Elevated antibody levels to ganglioside-monosialic acid (GM1), and the neutral glycolipid, asialo GM1 are associated with motor or sensorimotor neuropathies, particularly multifocal motor neuropathy. Anti-GM1 may occur as IgM (polyclonal or monoclonal) or IgG antibodies. These antibodies may also be found in patients with diverse connective tissue diseases as well as normal individuals. GD1a antibodies are associated with different variants of Guillain-Barre syndrome (GBS) particularly acute motor axonal neuropathy while GD1b antibodies are predominantly found in sensory ataxic neuropathy syndrome. Anti-GQ1b antibodies are seen in more than 80 percent of patients with Miller-Fisher syndrome and may be elevated in GBS patients with ophthalmoplegia. The role of isolated anti-GM2 antibodies is unknown. These tests by themselves are not diagnostic and should be used in conjunction with other clinical parameters to confirm disease.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

EER Motor Sensory Neuropathy Comp	See Note	
	Authorized individuals can access the ARUP Enhanced Report with an ARUP Connect account using the following link.	
	Your local lab can assist you in obtaining the patient report if you don't have a Connect account.	

Immunoglobulin G	1369 mg/dL	(Ref Interval: 768-1632)
Immunoglobulin A	270 mg/dL	(Ref Interval: 68-408)
Immunoglobulin M	46 mg/dL	(Ref Interval: 35-263)
Total Protein, Serum	7.6 g/dL	(Ref Interval: 6.3-8.2)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

Albumin	4.23 g/dL	(Ref Interval: 3.75-5.01)
Alpha 1 Globulin	0.32 g/dL	(Ref Interval: 0.19-0.46)
Alpha 2 Globulin	0.89 g/dL	(Ref Interval: 0.48-1.05)
Beta Globulin	0.99 g/dL	(Ref Interval: 0.48-1.10)
Gamma	1.18 g/dL	(Ref Interval: 0.62-1.51)
Monoclonal Protein	Not Applicable g/dL	(Ref Interval: <=0.00)
Immunofixation	IFE Done	
SPEP/IFE Interpretation	See Note Normal SPEP pattern. IFE gel shows a normal pattern; no monoclonal proteins seen.	

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
MAG Antibody, IgM Elisa	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SGPG Antibody, IgM	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Purkinje Cell/Neuronal Nuclear IgG Scrn	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Asialo-GM1 Antibodies, IgG/IgM	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GM1 Antibodies, IgG/IgM	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GD1a Antibodies, IgG/IgM	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GD1b Antibodies, IgG/IgM	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
GQ1b Antibodies, IgG/IgM	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
EER Motor Sensory Neuropathy Comp	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin G	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin A	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunoglobulin M	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Total Protein, Serum	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Albumin	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Alpha 1 Globulin	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Alpha 2 Globulin	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Beta Globulin	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Gamma	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Monoclonal Protein	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Immunofixation	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SPEP/IFE Interpretation	26-029-109008	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: