

Client: ARUP Example Report Only
500 Chipeta Way
Salt Lake City, UT 84108
UNITED STATES

Physician: ARUP, ARUP

**Patient: HPE PAN, NEGATIVE
EXAMPLE**

DOB

Sex: Female

Patient Identifiers: 41615

Visit Number (FIN): 41940

Collection Date: 8/16/2022 13:54

Holoprosencephaly Panel, Sequencing and Deletion/Duplication

ARUP test code 2008848

Holoprosencephaly Panel Specimen	whole Blood
Holoprosencephaly Panel Interp	<p>Negative</p> <p>INDICATION FOR TESTING Lobar holoprosencephaly</p> <p>RESULT No pathogenic variants were detected in any of the genes tested.</p> <p>INTERPRETATION No pathogenic variants were identified by massively parallel sequencing of the coding regions and exon-intron boundaries of the genes tested. No large exonic deletions and duplications were identified in the genes tested. This result decreases the likelihood of, but does not exclude, a diagnosis of a heritable form of holoprosencephaly. Please refer to the background information included in this report for a list of the genes analyzed and limitations of this test.</p> <p>RECOMMENDATIONS Medical screening and management should rely on clinical findings and family history. Genetic consultation is recommended.</p> <p>COMMENTS Benign and likely benign variants are not reported. Variants in the following region(s) may not be detected by NGS with sufficient confidence in this sample due to technical limitations; reportable variants are confirmed by Sanger sequencing: NONE</p> <p>This result has been reviewed and approved by [REDACTED]</p> <p>BACKGROUND INFORMATION: Holoprosencephaly Panel Sequencing and Deletion/Duplication</p> <p>CHARACTERISTICS: Holoprosencephaly (HPE) originates from failed midline delineation during early embryonic development and results in partial or complete failure of the prosencephalon to divide into hemispheres. Subtypes of HPE are based on the degree of brain separation and are typically classified as lobar, semilobar, alobar, and middle interhemispheric variant. Microform HPE is an additional subtype that has HPE-related craniofacial anomalies without structural brain defects. The range of clinical findings varies significantly within and between each subtype, including craniofacial characteristics, developmental delay, and neurological impacts. HPE may be an isolated finding or part of a broader syndrome.</p>

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: HPE PAN, NEGATIVE EXAMPLE
ARUP Accession: 22-228-114381
Patient Identifiers: 41615
Visit Number (FIN): 41940
Page 1 of 3 | Printed: 8/16/2022 2:13:32 PM

EPIDEMIOLOGY: The incidence is 1 in 10,000 live births.

CAUSE: 25-60 percent of cases are due to underlying cytogenetic abnormalities, including numerical or structural chromosome anomalies and pathogenic copy number variations. Pathogenic germline variants in single genes account for 25 percent of cases.

INHERITANCE: Dependent on etiology; autosomal dominant for genes tested in this panel

PENETRANCE: Incomplete with highly variable expression
CLINICAL SENSITIVITY: 25 percent

GENES TESTED: CDON; FGFR1*; GLI2; PTCH1; SHH; SIX3; TGIF1; ZIC2*
*One or more exons are not covered by sequencing and/or deletion/duplication analysis for the indicated gene; see limitations section below.

METHODOLOGY: Probe hybridization-based capture of all coding exons and exon-intron junctions of the targeted genes, followed by massively parallel sequencing. Sanger sequencing was performed as necessary to fill in regions of low coverage and to confirm reported variants that do not meet acceptable quality metrics. A proprietary bioinformatic algorithm was used to detect large (single exon-level or larger) deletions or duplications in the indicated genes. Large deletions/duplications confirmed using an orthogonal exon-level microarray. Human genome build 19 (Hg 19) was used for data analysis.

ANALYTICAL SENSITIVITY/SPECIFICITY: The analytical sensitivity is approximately 99 percent for single nucleotide variants (SNVs) and greater than 93 percent for insertions/duplications/deletions (indels) from 1-10 base pairs in size. Indels greater than 10 base pairs may be detected but the analytical sensitivity may be reduced. Deletions of 2 exons or larger are detected with sensitivity greater than 97 percent; single exon deletions are detected with 62 percent sensitivity. Duplications of 3 exons or larger are detected at greater than 83 percent sensitivity. Specificity is greater than 99.9 percent for all variant classes.

LIMITATIONS: A negative result does not exclude a heritable form of holoprosencephaly. This test only detects variants within the coding regions and intron-exon boundaries of the targeted genes. Deletions/duplications/insertions of any size may not be detected by massively parallel sequencing. Regulatory region variants and deep intronic variants will not be identified. Precise breakpoints for large deletions or duplications are not determined in this assay and single exon deletions/duplications may not be detected based on the breakpoints of the rearrangement. The actual breakpoints for the deletion or duplication may extend beyond or be within the exon(s) reported. This test is not intended to detect duplications of 2 or fewer exons in size, though these may be identified. Single exon deletions are reported but called at a lower sensitivity. Diagnostic errors can occur due to rare sequence variations. In some cases, variants may not be identified due to technical limitations caused by the presence of pseudogenes, repetitive, or homologous regions. This test is not intended to detect low-level mosaic or somatic variants, gene conversion events, complex inversions, translocations, mitochondrial DNA (mtDNA) mutations, or repeat expansions. Interpretation of this test result may be impacted if this patient has had an allogeneic stem cell transplantation. Noncoding transcripts were not analyzed.

The following regions are not sequenced due to technical limitations of the assay:

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: HPE PAN, NEGATIVE EXAMPLE
ARUP Accession: 22-228-114381
Patient Identifiers: 41615
Visit Number (FIN): 41940
Page 2 of 3 | Printed: 8/16/2022 2:13:32 PM

FGFR1 (NM_001354367) exon(s) 18
 FGFR1 (NM_001354369) exon(s) 18
 FGFR1 (NM_001354370) exon(s) 17
 ZIC2 (NM_007129) partial exon(s) 3(Chr13:100637736-100637843)
 Single exon deletions/duplications will not be called for the following exons: FGFR1 (NM_001354367, NM_001354369) 18; FGFR1 (NM_001354370) 17

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the U.S. Food and Drug Administration. This test was performed in a CLIA-certified laboratory and is intended for clinical purposes.

Counseling and informed consent are recommended for genetic testing. Consent forms are available online.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Holoprosencephaly Panel Specimen	22-228-114381	8/16/2022 1:54 00 PM	8/16/2022 1:55 02 PM	8/16/2022 2:10:00 PM
Holoprosencephaly Panel Interp	22-228-114381	8/16/2022 1:54 00 PM	8/16/2022 1:55 02 PM	8/16/2022 2:10:00 PM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: HPE PAN, NEGATIVE EXAMPLE
 ARUP Accession: 22-228-114381
 Patient Identifiers: 41615
 Visit Number (FIN): 41940
 Page 3 of 3 | Printed: 8/16/2022 2:13:32 PM