

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 2/13/1987
Gender: Male
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 01/01/2017 12:34

Herpes Simplex Virus Type 1 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050292

HSV 1 Glycoprotein G Ab, IgG

57.40 IV H (Ref Interval: <=0.89)

REFERENCE INTERVAL: HSV 1 Glycoprotein G Ab, IgG

- 0.89 IV or less Negative - No significant level of detectable IgG antibody to HSV type 1 glycoprotein G.
- 0.90 - 1.09 IV Equivocal - Questionable presence of IgG antibody to HSV type 1 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
- 1.10 IV or greater ... Positive - IgG antibody to HSV type 1 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

Herpes Simplex Virus Type 2 Glycoprotein G-Specific Antibody, IgG by CIA

ARUP test code 0050294

HSV 2 Glycoprotein G Antibody, IgG

0.71 IV (Ref Interval: <=0.89)

REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

- 0.89 IV or less Negative - No significant level of detectable IgG antibody to HSV type 2 glycoprotein G.
- 0.90 - 1.09 IV Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
- 1.10 IV or greater Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-129-109823
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 1 of 6 | Printed: 12/7/2020 2:09:01 PM
4848

Encephalitis Panel with Reflex to Herpes Simplex Virus Types 1 and 2 Glycoprotein G-Specific Antibodies, IgG, Serum

ARUP test code 2008915

HSV Type 1/2 Combined Ab, IgG

>22.40 IV

Specimen tested positive for Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG. ARUP test codes 0050292 and 0050294 will be added. Additional charges apply.

HSV Type 1 and Type 2 Glycoprotein G-Specific Antibodies, IgG to follow.

INTERPRETIVE INFORMATION: HSV 1/2 COMBINED Ab SCREEN, IgG
 0.89 IV or less.....Not Detected
 0.90-1.09 IV.....Indeterminate- Repeat testing
 in 10-14 days may be helpful.
 1.10 IV or greater.....Detected

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

HSV 1 and/or 2 Abs, IgM by ELISA

1.31 IV H (Ref Interval: <=0.89)

INTERPRETIVE INFORMATION: Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgM by ELISA

0.89 IV or Less Not Detected
 0.90 - 1.09 IV Indeterminate- Repeat testing in
 10-14 days may be helpful.
 1.10 IV or Greater Detected-IgM antibody to HSV
 detected, which may indicate a
 current or recent infection.
 However, low levels of IgM
 antibodies may occasionally
 persist for more than 12
 months post-infection.

West Nile Virus Ab, IgG, Ser

0.59 IV

(Ref Interval: <=1.29)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
 ARUP Accession: 20-129-109823
 Patient Identifiers: 01234567890ABCD, 012345
 Visit Number (FIN): 01234567890ABCD
 Page 2 of 6 | Printed: 12/7/2020 2:09:01 PM
 4848

INTERPRETIVE INFORMATION: West Nile Virus Ab, IgG by ELISA, Serum

- 1.29 IV or less Negative - No significant level of west Nile virus IgG antibody detected.
- 1.30 - 1.49 IV Equivocal - Questionable presence of West Nile virus IgG antibody detected. Repeat testing in 10-14 days may be helpful.
- 1.50 IV or greater Positive - Presence of IgG antibody to West Nile virus detected, suggestive of current or past infection.

This test is intended to be used as a semi-quantitative means of detecting West Nile virus-specific IgG in serum samples in which there is a clinical suspicion of West Nile virus infection. This test should not be used solely for quantitative purposes, nor should the results be used without correlation to clinical history or other data. Because other members of the Flaviviridae family, such as St. Louis encephalitis virus, show extensive cross-reactivity with West Nile virus, serologic testing specific for these species should be considered.

Seroconversion between acute and convalescent sera is considered strong evidence of current or recent infection. The best evidence for infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

West Nile Virus Ab, IgM, Ser

0.04 IV (Ref Interval: <=0.89)
INTERPRETIVE INFORMATION: West Nile Virus Ab, IgM by ELISA, Serum

- 0.89 IV or less Negative - No significant level of West Nile virus IgM antibody detected.
- 0.90-1.10 IV Equivocal - Questionable presence of West Nile virus IgM antibody detected. Repeat testing in 10-14 days may be helpful.
- 1.11 IV or greater ... Positive - Presence of IgM antibody to West Nile virus detected, suggestive of current or recent infection.

This test is intended to be used as a semi-quantitative means of detecting West Nile virus-specific IgM in serum samples in which there is a clinical suspicion of West Nile virus infection. This test should not be used solely for quantitative purposes, nor should the results be used without correlation to clinical history or other data. Because other members of the Flaviviridae family, such as St. Louis encephalitis virus, show extensive cross-reactivity with West Nile virus, serologic testing specific for these species should be considered.

Seroconversion between acute and convalescent sera is considered strong evidence of current or recent infection. The best evidence for infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Mumps Virus Antibody, IgG

<5.0 AU/mL

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-129-109823
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 3 of 6 | Printed: 12/7/2020 2:09:01 PM
4848

INTERPRETIVE INFORMATION: Mumps Ab, IgG by CIA

- 8.9 AU/mL or less Negative - No significant level of detectable IgG mumps virus antibody
- 9.0-10.9 AU/mL Equivocal - Repeat testing in 10-14 days may be helpful
- 11.0 AU/mL or greater: Positive - IgG antibody to mumps virus detected, which may indicate a current or past exposure/immunization to mumps virus.

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Mumps Virus Antibody, IgM

0.33 IV (Ref Interval: <=0.79)

INTERPRETIVE INFORMATION: Mumps Virus Antibody, IgM

- 0.79 IV or less: Negative - No significant level of detectable IgM antibody to mumps virus.
- 0.80 - 1.20 IV: Equivocal - Borderline levels of IgM antibody to mumps virus. Repeat testing in 10-14 days may be helpful.
- 1.21 IV or greater: Positive - Presence of IgM antibody to mumps virus detected, which may indicate a current or recent infection. However, low levels of IgM antibody may occasionally persist for more than 12 months post-infection or immunization.

Measles, Rubeola, Antibody IgG

83.0 AU/mL

INTERPRETIVE INFORMATION: Measles (Rubeola) Antibody, IgG

- 13.4 AU/mL or less..... Negative - No significant level of detectable measles (rubeola) IgG antibody.
- 13.5-16.4 AU/mL Equivocal - Repeat testing in 10-14 days may be helpful.
- 16.5 AU/mL or greater Positive - IgG antibody to measles (rubeola) detected which may indicate a current or past exposure/immunization to measles (rubeola).

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Measles, Rubeola, Antibody IgM

0.31 AU

(Ref Interval: 0.00-0.79)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

INTERPRETIVE INFORMATION: Measles (Rubeola) Antibody, IgM

0.79 AU or less Negative - No significant level of IgM antibody to measles (Rubeola) virus detected.
 0.80 - 1.20 AU Equivocal - Repeat testing in 10-14 days may be helpful.
 1.21 AU or greater Positive - IgM antibody to measles (Rubeola) virus detected. Suggestive of a current or recent infection or immunization. However, low levels of IgM antibodies may occasionally persist for more than 12 months post-infection or immunization.

Varicella-Zoster Virus Ab, IgG

3386.0 IV

INTERPRETIVE INFORMATION: VZV Ab, IgG

134.9 IV or less Negative - No significant level of detectable IgG varicella-zoster antibody.
 135.0 - 164.9 IV Equivocal - Repeat testing in 10-14 days may be helpful.
 165.0 IV or greater Positive - IgG antibody to varicella-zoster detected, which may indicate a current or past varicella-zoster infection.

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Varicella-Zoster Virus Antibody, IgM

0.20 ISR (Ref Interval: <=0.90)

INTERPRETIVE INFORMATION: Varicella-Zoster Virus Antibody, IgM

0.90 ISR or less Negative - No significant level of detectable varicella-zoster virus IgM antibody.
 0.91-1.09 ISR Equivocal - Repeat testing in 10-14 days may be helpful.
 1.10 ISR or greater Positive - Significant level of detectable varicella-zoster virus IgM antibody. Indicative of current or recent infection. However, low levels of IgM antibodies may occasionally persist for more than 12 months post-infection or immunization.

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
HSV Type 1/2 Combined Ab, IgG	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/9/2020 10:29:00 PM
HSV 1 and/or 2 Abs, IgM by ELISA	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/10/2020 7:57:00 AM
West Nile Virus Ab, IgG, Ser	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/13/2020 10:40:00 PM
West Nile Virus Ab, IgM, Ser	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/13/2020 10:40:00 PM
Mumps Virus Antibody, IgG	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/12/2020 2:48:00 PM
Mumps Virus Antibody, IgM	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/13/2020 7:11:00 AM
HSV 1 Glycoprotein G Ab, IgG	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 10:28:38 PM	5/10/2020 1:21:00 AM
HSV 2 Glycoprotein G Antibody, IgG	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 10:28:38 PM	5/10/2020 1:20:00 AM
Measles, Rubella, Antibody IgG	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/12/2020 2:47:00 PM
Measles, Rubella, Antibody IgM	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/13/2020 7:17:00 AM
Varicella-Zoster Virus Ab, IgG	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/12/2020 2:45:00 PM
Varicella-Zoster Virus Antibody, IgM	20-129-109823	5/8/2020 6:30:00 AM	5/9/2020 2:48:37 PM	5/14/2020 6:27:00 AM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-129-109823
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 6 of 6 | Printed: 12/7/2020 2:09:01 PM
4848