

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 10/27/1955  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Drug Profile, Screen with Reflex to Quantitation**

ARUP test code 2012312

Amphetamines, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 300)
Barbiturates, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 200)
Benzodiazepines, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 200)
Buprenorphine, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 5)
Carisoprodol, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 100)
Cocaine, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 150)
Creatinine, Urine	54.0 mg/dL	(Ref Interval: 20.0-400.0)
Ethyl Glucuronide Screen w/Rflx, Urine	Negative ng/mL	(Ref Interval: Cutoff 500) INTERPRETIVE INFORMATION: Ethyl Glucuronide Screen with Reflex to Confirmation, Urine  Ethyl glucuronide is a direct metabolite of ethanol and can be detected up to 80 hours in urine after ethanol ingestion. The cutoff for positive by immunoassay is set at 500 ng/mL. A positive result will be confirmed by liquid chromatography tandem mass spectrometry (LC-MS/MS).
Fentanyl, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 1)
Meperidine, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 200)

**H=High, L=Low, \*=Abnormal, C=Critical**

Methadone, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 150)
Opiates, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 300)
Oxycodone/Oxymorphone, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 100)
Phencyclidine, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 25)
Propoxyphene, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 300)
Tapentadol, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 200)
THC, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 20)
Tramadol, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 100)
Zolpidem, Urn, Screen	Negative ng/mL	(Ref Interval: Cutoff 20)

**Screen, Urine Interpretation**

**See Note**

**INTERPRETIVE INFORMATION:**

The absence of expected drug(s) and/or drug metabolite(s) may indicate non-compliance, inappropriate timing of specimen collection relative to drug administration, poor drug absorption, diluted/adulterated urine, or limitations of testing. The concentration at which the screening test can detect a drug or metabolite varies. Specimens for which drugs or drug classes are detected by the screen are reflexed to a second, more specific technology (GC/MS and/or LC-MS/MS). The concentration value must be greater than or equal to the cutoff to be reported as positive. Interpretive questions should be directed to the laboratory.

For medical purposes only; not valid for forensic use.

**H=High, L=Low, \*=Abnormal, C=Critical**

Unless otherwise indicated, testing performed at:

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Amphetamines, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Barbiturates, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Benzodiazepines, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Buprenorphine, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Carisoprodol, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Cocaine, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Creatinine, Urine	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Ethyl Glucuronide Screen w/Rflx, Urine	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Fentanyl, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Meperidine, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Methadone, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Opiates, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Oxycodone/Oxymorphone, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phencyclidine, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Propoxyphene, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Tapentadol, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
THC, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Tramadol, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Zolpidem, Urn, Screen	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Screen, Urine Interpretation	22-272-111595	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com  
500 Chipeta Way, Salt Lake City, UT 84108-1221  
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example  
ARUP Accession: 22-272-111595  
Patient Identifiers: 01234567890ABCD, 012345  
Visit Number (FIN): 01234567890ABCD  
Page 3 of 3 | Printed: 10/3/2022 2:42:45 PM  
4848