

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 4/17/1946
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 01/01/2017 12:34

Coxiella burnetii (Q-Fever) Antibodies, IgG and IgM, Phase I and II with Reflex to Titer

ARUP test code 2012634

C. Burnetii Abs, IgG Phase I Screen

Positive * (Ref Interval: Negative)

INTERPRETIVE INFORMATION: C. Burnetii Abs, IgG Phase I Screen

Acute Q fever is best demonstrated by a four-fold rise in phase II IgG titers when comparing two serum samples collected 3-6 weeks apart, and testing is performed in the same laboratory at the same time. Phase I IgG titers can increase during seroconversion. However, in the case of acute infection, the phase I titer should remain lower than the phase II titer. IgM antibodies to phase II antigens provide ancillary information to IgG titers. Phase II IgM titers develop in the same time period of phase II IgG titers and can persist for over a year. A single phase II IgM positive result on an acute sample represents an early conversion or a false positive; testing of a convalescent serum is necessary. In the absence of an acute sample, a single convalescent serum sample with a phase II IgG titer greater than 1:128 in a patient who has been ill longer than 1 week indicates probable acute Q fever. Chronic Q fever is best demonstrated by a phase I titer greater than the phase II IgG titer. Phase I IgM antibodies may also develop concurrently with phase I IgG antibodies. However, in the absence of a phase I IgG titer, the diagnostic value of a phase I IgM titer is limited. Phase I and phase II IgM and IgG titers may remain elevated for months or years after acute infection or during convalescence.

C. Burnetii Abs, IgG Phase II Screen

Positive * (Ref Interval: Negative)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

INTERPRETIVE INFORMATION: C. Burnetii Abs, IgG Phase II Screen

Acute Q fever is best demonstrated by a four-fold rise in phase II IgG titers when comparing two serum samples collected 3-6 weeks apart, and testing is performed in the same laboratory at the same time. Phase I IgG titers can increase during seroconversion. However, in the case of acute infection, the phase I titer should remain lower than the phase II titer. IgM antibodies to phase II antigens provide ancillary information to IgG titers. Phase II IgM titers develop in the same time period of phase II IgG titers and can persist for over a year. A single phase II IgM positive result on an acute sample represents an early conversion or a false positive; testing of a convalescent serum is necessary. In the absence of an acute sample, a single convalescent serum sample with a phase II IgG titer greater than 1:128 in a patient who has been ill longer than 1 week indicates probable acute Q fever. Chronic Q fever is best demonstrated by a phase I titer greater than the phase II IgG titer. Phase I IgM antibodies may also develop concurrently with phase I IgG antibodies. However, in the absence of a phase I IgG titer, the diagnostic value of a phase I IgM titer is limited. Phase I and phase II IgM and IgG titers may remain elevated for months or years after acute infection or during convalescence.

C. burnetii (Q-Fever) Ab, Phase I IgM

Positive * (Ref Interval: Negative)

INTERPRETIVE INFORMATION: C. burnetii (Q-Fever) Ab, Phase I IgM

Acute Q fever is best demonstrated by a four-fold rise in phase II IgG titers when comparing two serum samples collected 3-6 weeks apart, and testing is performed in the same laboratory at the same time. Phase I IgG titers can increase during seroconversion. However, in the case of acute infection, the phase I titer should remain lower than the phase II titer. IgM antibodies to phase II antigens provide ancillary information to IgG titers. Phase II IgM titers develop in the same time period of phase II IgG titers and can persist for over a year. A single phase II IgM positive result on an acute sample represents an early conversion or a false positive; testing of a convalescent serum is necessary. In the absence of an acute sample, a single convalescent serum sample with a phase II IgG titer greater than 1:128 in a patient who has been ill longer than 1 week indicates probable acute Q fever. Chronic Q fever is best demonstrated by a phase I titer greater than the phase II IgG titer. Phase I IgM antibodies may also develop concurrently with phase I IgG antibodies. However, in the absence of a phase I IgG titer, the diagnostic value of a phase I IgM titer is limited. Phase I and phase II IgM and IgG titers may remain elevated for months or years after acute infection or during convalescence.

C. burnetii (Q-Fever) Ab, Phase II IgM

Positive * (Ref Interval: Negative)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-167-104914
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 2 of 4 | Printed: 12/7/2020 2:53:15 PM
4848

INTERPRETIVE INFORMATION: C. burnetii (Q-Fever) Ab,
Phase II IgM

Acute Q fever is best demonstrated by a four-fold rise in phase II IgG titers when comparing two serum samples collected 3-6 weeks apart, and testing is performed in the same laboratory at the same time. Phase I IgG titers can increase during seroconversion. However, in the case of acute infection, the phase I titer should remain lower than the phase II titer. IgM antibodies to phase II antigens provide ancillary information to IgG titers. Phase II IgM titers develop in the same time period of phase II IgG titers and can persist for over a year. A single phase II IgM positive result on an acute sample represents an early conversion or a false positive; testing of a convalescent serum is necessary. In the absence of an acute sample, a single convalescent serum sample with a phase II IgG titer greater than 1:128 in a patient who has been ill longer than 1 week indicates probable acute Q fever. Chronic Q fever is best demonstrated by a phase I titer greater than the phase II IgG titer. Phase I IgM antibodies may also develop concurrently with phase I IgG antibodies. However, in the absence of a phase I IgG titer, the diagnostic value of a phase I IgM titer is limited. Phase I and phase II IgM and IgG titers may remain elevated for months or years after acute infection or during convalescence.

Coxiella burnetii (Q-Fever) Antibody, IgG Phase I Titer (Reflex for NEW TEST QFG1 SCRIN Not Orderable by Clients)

ARUP test code 2012630

Q-Fever Phase I IgG Titer	1:1024	(Ref Interval: < 1:16)
---------------------------	--------	------------------------

Coxiella burnetii (Q-Fever) Antibody, IgM Phase I Titer (Reflex for NEW TEST QFG1 SCRIN Not Orderable by Clients)

ARUP test code 2012643

Q-Fever Phase I IgM Titer	1:256	(Ref Interval: < 1:16)
---------------------------	-------	------------------------

Coxiella burnetii (Q-Fever) Antibody, IgG Phase II Titer (Reflex for NEW TEST QFG2 SCRIN Not Orderable by Clients)

ARUP test code 2012632

Q-Fever Phase II IgG Titer	1:65536	(Ref Interval: < 1:16)
----------------------------	---------	------------------------

Coxiella burnetii (Q-Fever) Antibody, IgM Phase II Titer (Reflex for NEW TEST QFG1 SCRIN Not Orderable by Clients)

ARUP test code 2012645

Q-Fever Phase II IgM Titer	1:1024	(Ref Interval: < 1:16)
----------------------------	--------	------------------------

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-167-104914
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 3 of 4 | Printed: 12/7/2020 2:53:15 PM
4848

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
C. Burnetii Abs, IgG Phase I Screen	20-167-104914	6/15/2020 7:49:00 AM	6/16/2020 2:24:01 AM	6/17/2020 9:35:00 PM
C. Burnetii Abs, IgG Phase II Screen	20-167-104914	6/15/2020 7:49:00 AM	6/16/2020 2:24:01 AM	6/17/2020 9:35:00 PM
C. burnetii (Q-Fever) Ab, Phase I IgM	20-167-104914	6/15/2020 7:49:00 AM	6/16/2020 2:24:01 AM	6/17/2020 9:35:00 PM
C. burnetii (Q-Fever) Ab, Phase II IgM	20-167-104914	6/15/2020 7:49:00 AM	6/16/2020 2:24:01 AM	6/17/2020 9:35:00 PM
Q-Fever Phase I IgG Titer	20-167-104914	6/15/2020 7:49:00 AM	6/17/2020 9:35:26 PM	6/17/2020 9:36:00 PM
Q-Fever Phase I IgM Titer	20-167-104914	6/15/2020 7:49:00 AM	6/17/2020 9:35:25 PM	6/17/2020 9:36:00 PM
Q-Fever Phase II IgG Titer	20-167-104914	6/15/2020 7:49:00 AM	6/17/2020 9:35:24 PM	6/17/2020 9:35:00 PM
Q-Fever Phase II IgM Titer	20-167-104914	6/15/2020 7:49:00 AM	6/17/2020 9:35:26 PM	6/17/2020 9:36:00 PM

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Tracy I. George, MD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 20-167-104914
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 4 of 4 | Printed: 12/7/2020 2:53:15 PM
4848