

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 1/12/1982
Gender: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Non-Criteria Antiphospholipid Syndrome (APS) (aPs, aPt, aPs/aPt) Antibodies Panel

ARUP test code 2012729

Prothrombin Antibody, IgG	1 Units (Ref Interval: 0-19) INTERPRETIVE INFORMATION: Prothrombin Antibody, IgG IgG antibodies to prothrombin may be a risk factor for either venous or arterial thrombosis in antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies. If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.
Phosphatidylserine Antibody IgG	0 GPS (Ref Interval: 0-15) INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgG IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies. Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE. If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.
Phosphatidylserine Antibody IgM	9 MPS (Ref Interval: 0-21)

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

<p>INTERPRETIVE INFORMATION: Phosphatidylserine Ab, IgM</p> <p>The presence of IgG and/or IgM antibodies to phosphatidylserine (aPS) may be associated with a positive test for anti-cardiolipin autoantibodies (aCL) and risk for obstetric antiphospholipid syndrome (APS). Strong clinical correlation is recommended in the absence of lupus anticoagulant, IgG and/or IgM cardiolipin and/or beta2 glycoprotein antibodies.</p> <p>Isolated presence of IgM or IgG antibodies to aPS may have questionable clinical significance for APS and/or SLE.</p> <p>If results are positive, repeat testing with two or more specimens drawn at least 12 weeks apart to demonstrate persistence of antibodies.</p> <p>Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.</p>	
<p>Phosphatidylserine and Prothrombin IgG</p>	<p>6 Units (Ref Interval: 0-30)</p> <p>INTERPRETIVE INFORMATION: Phosphatidylserine and Prothrombin IgG</p> <p>Elevated and persistent aPS/PT IgG antibody (with or without lupus anticoagulant activity) may serve as a risk marker of thrombotic events in patients with certain autoimmune diseases, including antiphospholipid syndrome (APS) and systemic lupus erythematosus (SLE).</p>
<p>Phosphatidylserine and Prothrombin IgM</p>	<p>16 Units (Ref Interval: 0-30)</p> <p>INTERPRETIVE INFORMATION: Phosphatidylserine and Prothrombin IgM</p> <p>Elevated and persistent aPS/PT IgM antibody (with or without lupus anticoagulant activity) may serve as a risk marker of thrombotic events in patients with certain autoimmune diseases, including antiphospholipid syndrome (APS) and systemic lupus erythematosus (SLE).</p>

H=High, L=Low, *=Abnormal, C=Critical



VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
Prothrombin Antibody, IgG	24-355-116701	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine Antibody IgG	24-355-116701	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine Antibody IgM	24-355-116701	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine and Prothrombin IgG	24-355-116701	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Phosphatidylserine and Prothrombin IgM	24-355-116701	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical