

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: Unknown

Gender: Unknown

Patient Identifiers: 01234567890ABCD, 012345

Visit Number (FIN): 01234567890ABCD

Collection Date: 00/00/0000 00:00

Apolipoprotein E (APOE) Genotyping, Alzheimer Disease Risk

ARUP test code 2013341

APOE Specimen whole Blood

APOE Alzheimer Disease Risk, Genotype **e2/e3 ***

Section 79-1 of New York State Civil Rights Law requires informed consent be obtained from patients (or their legal guardians) prior to pursuing genetic testing. These forms must be kept on file by the ordering physician. Consent forms for genetic testing are available at www.aruplab.com. Incidental findings are not reported unless clinically significant but are available upon request.

Indication for testing: Determine APOE genotype for the purpose of Alzheimer disease risk assessment.

Heterozygous APOE e2/e3: This genotype is associated with a decreased risk for Alzheimer disease (AD); however, the diagnosis of AD is primarily based on clinical evaluation, and APOE genotype alone is not sufficient to diagnose or exclude AD.

This result has been reviewed and approved by Yuan Ji, Ph.D.

H=High, L=Low, *=Abnormal, C=Critical

BACKGROUND INFORMATION: Apolipoprotein E (APOE) Genotyping, Alzheimer Disease Risk

Characteristics: Alzheimer disease (AD), the most common cause of dementia, is characterized by progressive cognitive decline including memory, problem-solving skills, multi-step tasks, planning, and changes in personality. A clinical diagnosis of probable AD can be made based on clinical signs and neuroimaging, and the diagnosis is confirmed postmortem based on neuropathologic findings. The e4 allele of the APOE gene has been widely demonstrated to be associated with increased risk of AD. In individuals with a clinical diagnosis of AD, the presence of the e4 allele increases the likelihood that the diagnosis is correct, but is not diagnostic alone. APOE genotyping is not recommended for predicting AD risk in asymptomatic individuals. Prevalence of APOE e4: Heterozygosity and homozygosity for the e4 allele is present in approximately 25 percent and 1-2 percent of the general population, respectively. Inheritance of APOE e4: Semi-dominant. Penetrance of APOE e4: Incomplete and influenced by age, gender, ethnicity, family history and environmental factors. The e4 allele is neither necessary nor sufficient for diagnosing AD; therefore, not all individuals with AD have the e4 allele and not all individuals with the e4 allele will develop AD. Cause: Multi-factorial.

Variants Tested: Two single nucleotide polymorphisms in the APOE gene at codons 130 (rs429358) and 176 (rs7412). The e3 allele (Cysteine at 130 and Arginine at 176) is the most common in the general population. The e4 allele (Arginine at 130 and 176) is associated with increased AD risk. The e2 allele (Cysteine at codons 130 and 176) may be associated with a lower risk for AD but homozygosity has been associated with increased risk for type III hyperlipoproteinemia.

Clinical Sensitivity: Approximately 30-60 percent of individuals diagnosed with AD carry at least one e4 allele. The e4/e4 genotype is found in approximately 13 percent of the AD population and 20 percent of the familial AD population.

Methodology: Polymerase chain reaction (PCR) and fluorescence monitoring using hybridization probes.

Analytical Sensitivity and Specificity: 99 percent.

Limitations: Only the APOE alleles e2, e3 and e4 will be detected; rare alleles are not detected by this test. Diagnostic errors can occur due to rare sequence variations.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement C: aruplab.com/CS

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
APOE Specimen	18-361-118722	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
APOE Alzheimer Disease Risk, Genotype	18-361-118722	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical