

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example** 

**DOB** 1/2/1945 **Gender:** Male

Patient Identifiers: 01234567890ABCD, 012345

**Visit Number (FIN):** 01234567890ABCD **Collection Date:** 00/00/0000 00:00

## Apolipoprotein E (APOE) Genotyping, Alzheimer Disease Risk

ARUP test code 2013341

APOE Specimen

Whole Blood

APOE Alzheimer Disease Risk, Genotype

e3/e3

Homozygous APOE e3/e3: This is the most common genotype in the general population; it is not associated with increased Alzheimer disease (AD) risk. However, the diagnosis of AD is primarily based on clinical evaluation and APOE genotype alone is not sufficient to diagnose or exclude AD.

This result has been reviewed and approved by

H=High, L=Low, \*=Abnormal, C=Critical

4848



BACKGROUND INFORMATION: Apolipoprotein E (APOE) Genotyping, Alzheimer Disease Risk

Characteristics: Alzheimer disease (AD), the most common cause of dementia, is characterized by progressive cognitive decline including memory, problem-solving skills, multi-step tasks, planning, and changes in personality. A clinical diagnosis of probable AD can be made based on clinical signs and neuroimaging, and the diagnosis is confirmed postmortem based on neuropathologic findings. The e4 allele of the APOE gene has been widely demonstrated to be associated with increased risk of AD. In individuals with a clinical diagnosis of AD, the presence of the e4 allele increases the likelihood that the diagnosis is correct, but is not diagnostic alone. APOE genotyping is not recommended for predicting AD risk in asymptomatic individuals. Prevalence of APOE e4: Heterozygosity and homozygosity for the e4 allele is present in approximately 25 percent and 1-2 percent of the general population, respectively.

Inheritance of APOE e4: Incomplete and influenced by age, gender, ethnicity, family history and environmental factors. The e4 allele is neither necessary nor sufficient for diagnosing AD; therefore, not all individuals with AD have the e4 allele and not all individuals with the e4 allele will develop AD.

Cause: Multi-factorial.

Variants Tested: Two single nucleotide polymorphisms in the APOE gene at codons 130 (rs429358) and 176 (rs7412). The e3 allele (Cysteine at 130 and Arginine at 176) is the most common in the general population. The e4 allele (Arginine at 130 and 176) is associated with increased AD risk. The e2 allele (Cysteine at codons 130 and 176) may be associated with increased risk for type III hyperlipoproteinemia.

Clinical Sensitivity: Approximately 30-60 percent of individuals diagnosed with AD carry at least one e4 allele. The e4/e4 genotype is found in approximately 13 percent of the AD Dopulation and 20 percent of the familial AD population. Methodology: Polymerase chain reaction (PCR) and fluorescence monitoring using hybridization probes.

Analytical Sensitivity and Specificity: 99 percent.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

Counseling and informed consent are recommended for genetic testing. Consent forms are available online.

H=High, L=Low, \*=Abnormal, C=Critical

4848



VERIFIED/REPORTED DATES				
Procedure	Accession	Collected	Received	Verified/Reported
APOE Specimen	22-088-402390	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
APOE Alzheimer Disease Risk, Genotype	22-088-402390	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Patient: Patient, Example ARUP Accession: 22-088-402390 Patient Identifiers: 01234567890ABCD, 012345 Visit Number (FIN): 01234567890ABCD Page 3 of 3 | Printed: 1/4/2023 3:00:29 PM

4848