

Client: Example Client ABC123 123 Test Drive Salt Lake City, UT 84108 UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

| 1/20/1975 |
|-------------------------|
| Male |
| 01234567890ABCD, 012345 |
| 01234567890ABCD |
| 00/00/0000 00:00 |
| |

Coccidioides Antibodies IgM by Immunoassay, CSF

| ARUP | test | code | 3000056 | |
|------|------|------|---------|--|
| | | | | |

| Coccidioides Antibody IgM ELISA, CSF | 4.5 IV | Н | (Ref Interval: <=0.2) | |
|--------------------------------------|--|------------|--|--|
| | Positive EIA results should be confirmed with immunodiffusion (ID) or the complement fixation (CF) test. When the ID or CF tests fail to corroborate the EIA results, the diagnosis is less firmly established. | | | |
| | INTERPRETIV | E INFORMAT | TION: Coccidioides Antibody IgM ELISA, CSF | |
| | 0.2 IV or | less: | Negative - No significant level of Coccidioides IgM antibody detected. | |
| | 0.3 IV or | greater: | Positive - Presence of IgM antibody to Coccidioides detected, suggestive of current or past infection. | |
| | In most symptomatic patients, IgM antibodies usually ap the second week of infection and disappear by the fourt Both tube precipitin (TP) and CF antigens are represent ELISA tests. | | | |

| VERIFIED/REPORTED DATES | | | | | | |
|--------------------------------------|---------------|------------------|------------------|-------------------|--|--|
| Procedure | Accession | Collected | Received | Verified/Reported | | |
| Coccidioides Antibody IgM ELISA, CSF | 22-337-113087 | 00/00/0000 00:00 | 00/00/0000 00:00 | 00/00/0000 00:00 | | |

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: