

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 5/10/1968  
**Gender:** Female  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**ADAMTS13 Inhibitor**

ARUP test code 3000228

ADAMTS13 Inhibitor

<0.4 BU (Ref Interval: <=0.4)

INTERPRETIVE INFORMATION: ADAMTS13 Inhibitor

The majority of cases of idiopathic thrombotic thrombocytopenic purpura (TTP) are caused by ADAMTS13 autoantibodies. Autoantibodies that neutralize ADAMTS13 function are found in approximately two-thirds of idiopathic cases and can be identified and titered by the ADAMTS13 inhibitor test. Non-neutralizing autoantibodies that result in increased ADAMTS13 clearance, but do not inhibit function, are found in approximately one-third of idiopathic TTP cases. Non-neutralizing antibodies are not detected in the inhibitor test but can be detected by ELISA (ADAMTS13 antibody test). ADAMTS13 autoantibodies are not present in congenital TTP (Upshaw-Schulman syndrome). Correlation with clinical information, ADAMTS13 activity, and other relevant laboratory testing is suggested.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
ADAMTS13 Inhibitor	22-087-119848	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical

Unless otherwise indicated, testing performed at: