

Client: Example Client ABC123  
123 Test Drive  
Salt Lake City, UT 84108  
UNITED STATES

Physician: Doctor, Example

**Patient: Patient, Example**

**DOB:** 7/4/1963  
**Gender:** Male  
**Patient Identifiers:** 01234567890ABCD, 012345  
**Visit Number (FIN):** 01234567890ABCD  
**Collection Date:** 00/00/0000 00:00

**Blastomyces Antigen Quantitative by EIA, Urine (Temporary Referral as of 08/19/19)**

ARUP test code 3001798

Blastomyces dermatitidis EIA, Interp

**Detected \* (Ref Interval: Not Detected)**

INTERPRETIVE INFORMATION: Blastomyces Ag Quant by ELISA, Urine

Less than 1.25 ng/ml = Not Detected

1.25-200 ng/mL = Detected

Greater than 200 ng/mL = Detected (above the limit of quantification)

The quantitative range of this assay is 1.25-200 ng/mL. Antigen concentrations greater than 200 ng/mL fall outside the linear range of the assay and cannot be accurately quantified.

This EIA test should be used in conjunction with other diagnostic procedures, including microbiological culture, histological examination of biopsy samples, and/or radiographic evidence, to aid in the diagnosis of blastomycosis.

Cross-reactivity with other endemic mycoses (Histoplasma and Coccidioides) may occur. Positive test results should be correlated with other clinical findings and relevant exposure history

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

Blastomyces Antigen, Urine

50.00 ng/mL

H=High, L=Low, \*=Abnormal, C=Critical

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Blastomyces dermatitidis EIA, Interp	19-232-105138	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Blastomyces Antigen, Urine	19-232-105138	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, \*=Abnormal, C=Critical