

Client: Example Client ABC123
123 Test Drive
Salt Lake City, UT 84108
UNITED STATES

Physician: Doctor, Example

Patient: Patient, Example

DOB: 4/10/1961
Sex: Female
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Collection Date: 00/00/0000 00:00

Connective Tissue Disease First Line Panel with Reflex

ARUP test code 3002463

Double-Stranded DNA (dsDNA) Ab IgG ELISA 10 IU (Ref Interval: 0-24)
 INTERPRETIVE INFORMATION: Double-Stranded DNA (dsDNA) Ab IgG ELISA
 24 IU or less.....Negative
 25-30 IU.....Borderline Positive
 30-60 IU.....Low Positive
 60-200 IU.....Positive
 201 IU or greater....Strong Positive

Positivity for anti-double stranded DNA (anti-dsDNA) IgG antibody is a diagnostic criterion of systemic lupus erythematosus (SLE). Specimens are initially screened by enzyme-linked immunosorbent assay (ELISA). If ordered as reflex (0050215), positive ELISA results (>24 IU) will be reflexed to a highly specific IFA titer (Crithidia luciliae indirect fluorescent test [CLIFT]) for confirmation. Some patients with early or inactive SLE may be positive for anti-dsDNA IgG by ELISA but negative by CLIFT. If the patient is negative by CLIFT but positive by ELISA and clinical suspicion remains, consider antinuclear antibody (ANA) testing by IFA. Additional information and recommendations for testing may be found at <https://arupconsult.com/content/systemic-lupus-erythematosus>.

Smith (ENA) Antibody, IgG 1 AU/mL (Ref Interval: 0-40)
 INTERPRETIVE INFORMATION: Smith (ENA) Antibody, IgG
 29 AU/mL or Less Negative
 30 - 40 AU/mL Equivocal
 41 AU/mL or Greater Positive

Smith antibody is highly specific (greater than 90 percent) for systemic lupus erythematosus (SLE) but only occurs in 30-35 percent of SLE cases. The presence of antibodies to Smith has variable associations with SLE clinical manifestations.

SSA-52 (Ro52) (ENA) Antibody, IgG 0 AU/mL (Ref Interval: 0-40)
 INTERPRETIVE INFORMATION: SSA-52 (Ro52) (ENA) Antibody, IgG
 29 AU/mL or Less Negative
 30 - 40 AU/mL Equivocal
 41 AU/mL or Greater Positive

SSA-52 (Ro52) and/or SSA-60 (Ro60) antibodies are associated with a diagnosis of Sjogren syndrome, systemic lupus erythematosus (SLE), and systemic sclerosis. SSA-52 antibody

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 25-212-401597
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 1 of 3 | Printed: 8/19/2025 11:44:33 AM

overlaps significantly with the major SSC-related antibodies. SSA-52 (Ro52) antibody occurs frequently in patients with inflammatory myopathies, often in the presence of interstitial lung disease.

SSA-60 (Ro60) (ENA) Antibody, IgG

0 AU/mL (Ref Interval: 0-40)
REFERENCE INTERVAL: SSA-60 (Ro60) (ENA) Antibody, IgG
29 AU/mL or Less Negative
30 - 40 AU/mL Equivocal
41 AU/mL or Greater Positive

Smith/RNP (ENA) Ab, IgG

2 Units (Ref Interval: 0-19)
INTERPRETIVE INFORMATION: Smith/RNP (ENA) Antibody, IgG
19 Units or Less Negative
20 to 39 Units Weak Positive
40 to 80 Units Moderate Positive
81 Units or greater Strong Positive

Smith/RNP antibodies are frequently seen in patients with mixed connective tissue disease (MCTD) and are also associated with other systemic autoimmune rheumatic diseases (SARDs) such as systemic lupus erythematosus (SLE), systemic sclerosis, and myositis. Antibodies targeting the Smith/RNP antigenic complex also recognize Smith antigens, therefore, the Smith antibody response must be considered when interpreting these results.

Jo-1 (Histidyl-tRNA Synthetase) Ab, IgG

0 AU/mL (Ref Interval: 0-40)
INTERPRETIVE INFORMATION: Jo-1 Antibody, IgG
29 AU/mL or less.....Negative
30-40 AU/mL.....Equivocal
41 AU/mL or greater.....Positive

Presence of Jo-1 (antihistidyl transfer RNA [t-RNA] synthetase) antibody is associated with polymyositis and may also be seen in patients with dermatomyositis. Jo-1 antibody is associated with pulmonary involvement (interstitial lung disease), Raynaud phenomenon, arthritis, and mechanic's hands (implicated in antisynthetase syndrome).

SSB (La) (ENA) Antibody, IgG

1 AU/mL (Ref Interval: 0-40)
INTERPRETIVE INFORMATION: SSB (La) (ENA) Ab, IgG
29 AU/mL or Less Negative
30 - 40 AU/mL Equivocal
41 AU/mL or Greater Positive

SSB (La) antibody is seen in 50-60% of Sjogren syndrome cases and is specific if it is the only ENA antibody present. 15-25% of patients with systemic lupus erythematosus (SLE) and 5-10% of patients with progressive systemic sclerosis (PSS) also have this antibody.

Scleroderma (Scl-70) (ENA) Antibody, IgG

0 AU/mL (Ref Interval: 0-40)
INTERPRETIVE INFORMATION: Scleroderma (Scl-70) (ENA) Ab, IgG
29 AU/mL or Less Negative
30 - 40 AU/mL Equivocal

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

41 AU/mL or Greater Positive

The presence of Scl-70 antibodies (also referred to as topoisomerase I, topo-I or ATA) is considered diagnostic for systemic sclerosis (SSc). Scl-70 antibodies alone are detected in about 20 percent of SSc patients and are associated with the diffuse form of the disease, which may include specific organ involvement and poor prognosis. Scl-70 antibodies have also been reported in a varying percentage of patients with systemic lupus erythematosus (SLE). Scl-70 (topo-1) is a DNA binding protein and anti-DNA/DNA complexes in the sera of SLE patients may bind to topo-I, leading to a false-positive result. The presence of Scl-70 antibody in sera may also be due to contamination of recombinant Scl-70 with DNA derived from cellular material used in immunoassays. Strong clinical correlation is recommended if both Scl-70 and dsDNA antibodies are detected.

Negative results do not necessarily rule out the presence of SSc. If clinical suspicion remains, consider further testing for centromere, RNA polymerase III and U3-RNP, PM/Scl, or Th/To antibodies.

VERIFIED/REPORTED DATES

Procedure	Accession	Collected	Received	Verified/Reported
Double-Stranded DNA (dsDNA) Ab IgG ELISA	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Smith (ENA) Antibody, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SSA-52 (Ro52) (ENA) Antibody, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SSA-60 (Ro60) (ENA) Antibody, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Smith/RNP (ENA) Ab, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Jo-1 (Histidyl-tRNA Synthetase) Ab, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
SSB (La) (ENA) Antibody, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00
Scleroderma (Scl-70) (ENA) Antibody, IgG	25-212-401597	00/00/0000 00:00	00/00/0000 00:00	00/00/0000 00:00

END OF CHART

H=High, L=Low, *=Abnormal, C=Critical

Unless otherwise indicated, testing performed at:

ARUP LABORATORIES | 800-522-2787 | aruplab.com
500 Chipeta Way, Salt Lake City, UT 84108-1221
Jonathan R. Genzen, MD, PhD, Laboratory Director

Patient: Patient, Example
ARUP Accession: 25-212-401597
Patient Identifiers: 01234567890ABCD, 012345
Visit Number (FIN): 01234567890ABCD
Page 3 of 3 | Printed: 8/19/2025 11:44:33 AM